

SCHRIFTENREIHE DER FAKULTÄT FÜR MATHEMATIK

A Compactness Result for the div-curl System with
Inhomogeneous Mixed Boundary Conditions
for Bounded Lipschitz Domains
and Some Applications

by

Dirk Pauly & Nathanael Skrepek

SM-UDE-825

2021

Received: March 11, 2021

**A COMPACTNESS RESULT FOR THE DIV-CURL SYSTEM WITH
INHOMOGENEOUS MIXED BOUNDARY CONDITIONS
FOR BOUNDED LIPSCHITZ DOMAINS
AND SOME APPLICATIONS**

DIRK PAULY AND NATHANAEL SKREPEK

ABSTRACT. For a bounded Lipschitz domain with Lipschitz interface we show the following *compactness theorem*: Any L^2 -bounded sequence of vector fields with L^2 -bounded rotations and L^2 -bounded divergences as well as L^2 -bounded tangential traces on one part of the boundary and L^2 -bounded normal traces on the other part of the boundary, contains a strongly L^2 -convergent subsequence. This generalises recent results for homogeneous mixed boundary conditions in [2, 4]. As applications we present a related *Friedrichs/Poincaré type estimate*, a *div-curl lemma*, and show that the Maxwell operator with mixed tangential and impedance boundary conditions (Robin type boundary conditions) has *compact resolvents*.

CONTENTS

| | | |
|------|---|---|
| 1. | Introduction | 1 |
| 2. | Notations | 2 |
| 3. | Preliminaries | 3 |
| 4. | Compact Embeddings | 4 |
| 5. | Applications | 5 |
| 5.1. | Friedrichs/Poincaré Type Estimates | 5 |
| 5.2. | A div-curl Lemma | 6 |
| 5.3. | Maxwell's Equations with Mixed Impedance Type Boundary Conditions | 6 |
| 5.4. | Wave Equation with Mixed Impedance Type Boundary Conditions | 8 |
| | References | 9 |

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^3$ be open with boundary Γ , composed of the boundary parts Γ_0 (tangential) and Γ_1 (normal). In [2, Theorem 4.7] the following version of Weck's selection theorem has been shown:

Theorem 1.1 (compact embedding for vector fields with homogeneous mixed boundary conditions). *Let (Ω, Γ_0) be a bounded strong Lipschitz pair and let ε be admissible. Then*

$$H_{\Gamma_0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1} H_{\Gamma_1}(\operatorname{div}, \Omega) \xrightarrow{\text{cpt}} L^2(\Omega).$$

Here, $\xrightarrow{\text{cpt}}$ denotes a compact embedding, and $-$ in classical terms and in the smooth case $-$ we have for a vector field E (n denotes the exterior unit normal at Γ)

$$E \in H_{\Gamma_0}(\operatorname{curl}, \Omega) \quad \Leftrightarrow \quad E \in L^2(\Omega), \quad \operatorname{curl} E \in L^2(\Omega), \quad n \times E|_{\Gamma_0} = 0,$$

Date: March 13, 2021.

Key words and phrases. compact embeddings, div-curl system, mixed boundary conditions, inhomogeneous boundary conditions.

We gratefully acknowledge ISem23 (23rd Internet Seminar 2019/2020, Evolutionary Equations, <http://www.mat.tuhh.de/isem23>) for providing the platform to start this research. The second author has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 765579.

$$E \in \varepsilon^{-1}H_{\Gamma_1}(\operatorname{div}, \Omega) \quad \Leftrightarrow \quad \varepsilon E \in L^2(\Omega), \quad \operatorname{div} \varepsilon E \in L^2(\Omega), \quad n \cdot \varepsilon E|_{\Gamma_1} = 0.$$

Note that Theorem 1.1 even holds for bounded weak Lipschitz pairs (Ω, Γ_0) . For exact definitions and notations see Section 2, and for a history of related compact embedding results see, e.g., [24, 19, 23, 5, 26, 7, 21] and [9]. The general importance of compact embeddings in a functional analytical setting (FA-ToolBox) for Hilbert complexes (such as de Rham, elasticity, biharmonic) is described, e.g., in [13, 15, 16, 14] and [17, 18, 1].

In this paper, we shall generalise Theorem 1.1 to the case of inhomogeneous boundary conditions, i.e., we will show that the compact embedding in Theorem 1.1 still holds if the space

$$H_{\Gamma_0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1}H_{\Gamma_1}(\operatorname{div}, \Omega)$$

is replaced by

$$\widehat{H}_{\Gamma_0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1}\widehat{H}_{\Gamma_1}(\operatorname{div}, \Omega),$$

where in classical terms and in the smooth case

$$\begin{aligned} E \in \widehat{H}_{\Gamma_0}(\operatorname{curl}, \Omega) &\quad \Leftrightarrow \quad E \in L^2(\Omega), \quad \operatorname{curl} E \in L^2(\Omega), \quad n \times E|_{\Gamma_0} \in L^2(\Gamma_0), \\ E \in \varepsilon^{-1}\widehat{H}_{\Gamma_1}(\operatorname{div}, \Omega) &\quad \Leftrightarrow \quad \varepsilon E \in L^2(\Omega), \quad \operatorname{div} \varepsilon E \in L^2(\Omega), \quad n \cdot \varepsilon E|_{\Gamma_1} \in L^2(\Gamma_1). \end{aligned}$$

The main result (compact embedding) is formulated in Theorem 4.1. As applications we show in Theorem 5.1 that the compact embedding implies a related Friedrichs/Poincaré type estimate, showing well-posedness of related systems of partial differential equations. Moreover, in Theorem 5.3 we prove that Theorem 4.1 yields a div-curl lemma. Note that corresponding results for exterior domains are straight forward using weighted Sobolev spaces, see [11, 12]. Another application is presented in Section 5.3 where we show that our compact embedding result implies compact resolvents of the Maxwell operator with inhomogeneous mixed boundary conditions, even of impedance type. We finally note in Section 5.4 that the corresponding result holds (in the simpler situation) for the impedance wave equation (acoustics) as well.

2. NOTATIONS

Throughout this paper, let $\Omega \subset \mathbb{R}^3$ be an open and bounded strong Lipschitz domain, and let ε be an *admissible* tensor (matrix) field, i.e., a symmetric, L^∞ -bounded, and uniformly positive definite tensor field $\varepsilon : \Omega \rightarrow \mathbb{R}^{3 \times 3}$. Moreover, let the boundary Γ of Ω be decomposed into two relatively open and strong Lipschitz subsets Γ_0 and $\Gamma_1 := \Gamma \setminus \overline{\Gamma_0}$ forming the interface $\overline{\Gamma_0} \cap \overline{\Gamma_1}$ for the mixed boundary conditions. See [2, 3, 4] for exact definitions. We call (Ω, Γ_0) a bounded strong Lipschitz pair.

The usual Lebesgue and Sobolev Hilbert spaces (of scalar or vector valued fields) are denoted by $L^2(\Omega)$, $H^1(\Omega)$, $H(\operatorname{curl}, \Omega)$, $H(\operatorname{div}, \Omega)$, and by $H_0(\operatorname{curl}, \Omega)$ and $H_0(\operatorname{div}, \Omega)$ we indicate the spaces with vanishing curl and div, respectively. Homogeneous boundary conditions are introduced in the strong sense as closures of respective test fields from

$$C_{\Gamma_0}^\infty(\Omega) := \{ \phi|_\Omega : \phi \in C^\infty(\mathbb{R}^3), \operatorname{supp} \phi \text{ compact, } \operatorname{dist}(\operatorname{supp} \phi, \Gamma_0) > 0 \},$$

i.e.,

$$H_{\Gamma_0}^1(\Omega) := \overline{C_{\Gamma_0}^\infty(\Omega)}^{H^1(\Omega)}, \quad H_{\Gamma_0}(\operatorname{curl}, \Omega) := \overline{C_{\Gamma_0}^\infty(\Omega)}^{H(\operatorname{curl}, \Omega)}, \quad H_{\Gamma_0}(\operatorname{div}, \Omega) := \overline{C_{\Gamma_0}^\infty(\Omega)}^{H(\operatorname{div}, \Omega)},$$

and we set $H_\emptyset^1(\Omega) := H^1(\Omega)$, $H_\emptyset(\operatorname{curl}, \Omega) := H(\operatorname{curl}, \Omega)$, and $H_\emptyset(\operatorname{div}, \Omega) := H(\operatorname{div}, \Omega)$. Spaces with vanishing curl and div are again denoted by $H_{\Gamma_0,0}(\operatorname{curl}, \Omega)$ and $H_{\Gamma_0,0}(\operatorname{div}, \Omega)$, respectively. Moreover, we introduce the cohomology space of Dirichlet/Neumann fields (generalised harmonic fields)

$$\mathcal{H}_{\Gamma_0, \Gamma_1, \varepsilon}(\Omega) := H_{\Gamma_0,0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1}H_{\Gamma_1,0}(\operatorname{div}, \Omega).$$

The $L^2(\Omega)$ -inner product and norm (of scalar or vector valued $L^2(\Omega)$ -spaces) will be denoted by $\langle \cdot, \cdot \rangle_{L^2(\Omega)}$ and $\| \cdot \|_{L^2(\Omega)}$, respectively, and the weighted Lebesgue space $L_\varepsilon^2(\Omega)$ is defined as $L^2(\Omega)$ (of vector fields) but being equipped with the weighted $L^2(\Omega)$ -inner product and norm $\langle \cdot, \cdot \rangle_{L_\varepsilon^2(\Omega)} := \langle \varepsilon \cdot, \cdot \rangle_{L^2(\Omega)}$ and $\| \cdot \|_{L_\varepsilon^2(\Omega)}$, respectively. The norms in, e.g., $H^1(\Omega)$ and $H(\operatorname{curl}, \Omega)$ are

denoted by $\|\cdot\|_{\mathbf{H}^1(\Omega)}$ and $\|\cdot\|_{\mathbf{H}(\text{curl},\Omega)}$, respectively. Orthogonality and orthogonal sum in $\mathbf{L}^2(\Omega)$ and $\mathbf{L}_\varepsilon^2(\Omega)$ are indicated by $\perp_{\mathbf{L}^2(\Omega)}$, $\perp_{\mathbf{L}_\varepsilon^2(\Omega)}$, and $\oplus_{\mathbf{L}^2(\Omega)}$, $\oplus_{\mathbf{L}_\varepsilon^2(\Omega)}$, respectively.

Finally, we introduce inhomogeneous tangential and normal \mathbf{L}^2 -boundary conditions in

$$\begin{aligned}\widehat{\mathbf{H}}_{\Gamma_0}(\text{curl}, \Omega) &:= \{E \in \mathbf{H}(\text{curl}, \Omega) : \tau_{\Gamma_0} E \in \mathbf{L}^2(\Gamma_0)\}, \\ \widehat{\mathbf{H}}_{\Gamma_1}(\text{div}, \Omega) &:= \{E \in \mathbf{H}(\text{div}, \Omega) : \nu_{\Gamma_1} E \in \mathbf{L}^2(\Gamma_1)\}\end{aligned}$$

with norms given by, e.g., $\|E\|_{\widehat{\mathbf{H}}_{\Gamma_0}(\text{curl}, \Omega)}^2 := \|E\|_{\mathbf{H}(\text{curl}, \Omega)}^2 + \|\tau_{\Gamma_0} E\|_{\mathbf{L}^2(\Gamma_0)}^2$. The definitions of the latter Hilbert spaces need some explanations:

Definition 2.1 (\mathbf{L}^2 -traces).

- (i) *The tangential trace of a vector field $E \in \mathbf{H}(\text{curl}, \Omega)$ is a well-defined tangential vector field $\tau_\Gamma E \in \mathbf{H}^{-1/2}(\Gamma)$ generalising the classical tangential trace $\tau_\Gamma \widetilde{E} = -n \times n \times \widetilde{E}|_\Gamma$ for smooth vector fields \widetilde{E} . By the notation $\tau_{\Gamma_0} E \in \mathbf{L}^2(\Gamma_0)$ we mean, that there exists a tangential vector field $E_{\Gamma_0} \in \mathbf{L}^2(\Gamma_0)$, such that for all vector fields $\Phi \in \mathbf{H}_{\Gamma_1}^1(\Omega)$ it holds*

$$\langle \text{curl } \Phi, E \rangle_{\mathbf{L}^2(\Omega)} - \langle \Phi, \text{curl } E \rangle_{\mathbf{L}^2(\Omega)} = \langle \tau_{\Gamma_0}^\times \Phi, E_{\Gamma_0} \rangle_{\mathbf{L}^2(\Gamma_0)}.$$

Then we set $\tau_{\Gamma_0} E := E_{\Gamma_0} \in \mathbf{L}^2(\Gamma_0)$. Here and in the following, the twisted tangential trace of the smooth vector field Φ is given by the tangential vector field $\tau_\Gamma^\times \Phi = n \times \Phi|_\Gamma \in \mathbf{L}^2(\Gamma)$ with $\tau_{\Gamma_1}^\times \Phi = \tau_\Gamma^\times \Phi|_{\Gamma_1} = 0$ and $\tau_{\Gamma_0}^\times \Phi = \tau_\Gamma^\times \Phi|_{\Gamma_0} \in \mathbf{L}^2(\Gamma_0)$. Note that $\tau_{\Gamma_0} E$ is well defined as $\tau_{\Gamma_0}^\times \mathbf{H}_{\Gamma_1}^1(\Omega)$ is dense in $\mathbf{L}_t^2(\Gamma_0) = \{v \in \mathbf{L}^2(\Gamma_0) : n \cdot v = 0\}$.

- (ii) *Analogously, the normal trace of a vector field $E \in \mathbf{H}(\text{div}, \Omega)$ is a well-defined function $\nu_\Gamma E \in \mathbf{H}^{-1/2}(\Gamma)$ generalising the classical normal trace $\nu_\Gamma \widetilde{E} = n \cdot \widetilde{E}|_\Gamma$ for smooth vector fields \widetilde{E} . Again, by the notation $\nu_{\Gamma_1} E \in \mathbf{L}^2(\Gamma_1)$ we mean, that for all functions $\phi \in \mathbf{H}_{\Gamma_0}^1(\Omega)$ it holds*

$$\langle \nabla \phi, E \rangle_{\mathbf{L}^2(\Omega)} + \langle \phi, \text{div } E \rangle_{\mathbf{L}^2(\Omega)} = \langle \sigma_{\Gamma_1} \phi, \nu_{\Gamma_1} E \rangle_{\mathbf{L}^2(\Gamma_1)}.$$

Here, the well-known scalar trace of the smooth function ϕ is given by $\sigma_\Gamma \phi = \phi|_\Gamma \in \mathbf{L}^2(\Gamma)$ with $\sigma_{\Gamma_0} \phi = \sigma_\Gamma \phi|_{\Gamma_0} = 0$ and $\sigma_{\Gamma_1} \phi = \sigma_\Gamma \phi|_{\Gamma_1} \in \mathbf{L}^2(\Gamma_1)$. Note that $\nu_{\Gamma_1} E$ is well defined as $\sigma_{\Gamma_1} \mathbf{H}_{\Gamma_0}^1(\Omega)$ is dense in $\mathbf{L}^2(\Gamma_1)$.

Remark 2.2 (\mathbf{L}^2 -traces). *Analogously to Definition 2.1 (i) and as*

$$\tau_{\Gamma_0}^\times \widetilde{E} \cdot \tau_{\Gamma_0} \widetilde{H} = (n \times \widetilde{E}) \cdot (-n \times n \times \widetilde{H}) = (n \times n \times \widetilde{E}) \cdot (n \times \widetilde{H}) = -\tau_{\Gamma_0} \widetilde{E} \cdot \tau_{\Gamma_0}^\times \widetilde{H}$$

holds on Γ_0 for smooth vector fields $\widetilde{E}, \widetilde{H}$, we can define the twisted tangential trace $\tau_{\Gamma_0}^\times E \in \mathbf{L}^2(\Gamma_0)$ of a vector field $E \in \mathbf{H}(\text{curl}, \Omega)$ as well by

$$\langle \text{curl } \Phi, E \rangle_{\mathbf{L}^2(\Omega)} - \langle \Phi, \text{curl } E \rangle_{\mathbf{L}^2(\Omega)} = -\langle \tau_{\Gamma_0} \Phi, \tau_{\Gamma_0}^\times E \rangle_{\mathbf{L}^2(\Gamma_0)}$$

for all vector fields $\Phi \in \mathbf{H}_{\Gamma_1}^1(\Omega)$.

3. PRELIMINARIES

In [4, Theorem 5.5], see [3, Theorem 7.4] for more details and compare to [2], the following theorem about the existence of regular potentials for the rotation with homogeneous mixed boundary conditions has been shown.

Theorem 3.1 (regular potential for curl with homogeneous mixed boundary conditions).

$$\mathbf{H}_{\Gamma_1,0}(\text{div}, \Omega) \cap \mathcal{H}_{\Gamma_0,\Gamma_1}(\Omega)^{\perp_{\mathbf{L}^2(\Omega)}} = \text{curl } \mathbf{H}_{\Gamma_1}(\text{curl}, \Omega) = \text{curl } \mathbf{H}_{\Gamma_1}^1(\Omega)$$

holds together with a regular potential operator mapping $\text{curl } \mathbf{H}_{\Gamma_1}(\text{curl}, \Omega)$ to $\mathbf{H}_{\Gamma_1}^1(\Omega)$ continuously. In particular, the latter ranges are closed subspaces of $\mathbf{L}^2(\Omega)$.

Moreover, we need [4, Theorem 5.2]:

Theorem 3.2 (Helmholtz decompositions with homogeneous mixed boundary conditions). *The ranges $\nabla \mathbf{H}_{\Gamma_0}^1(\Omega)$ and $\text{curl} \mathbf{H}_{\Gamma_1}(\text{curl}, \Omega)$ are closed subspaces of $\mathbf{L}^2(\Omega)$, and the $\mathbf{L}_\varepsilon^2(\Omega)$ -orthogonal Helmholtz decompositions*

$$\begin{aligned} \mathbf{L}_\varepsilon^2(\Omega) &= \nabla \mathbf{H}_{\Gamma_0}^1(\Omega) \oplus_{\mathbf{L}_\varepsilon^2(\Omega)} \varepsilon^{-1} \mathbf{H}_{\Gamma_1,0}(\text{div}, \Omega) \\ &= \mathbf{H}_{\Gamma_0,0}(\text{curl}, \Omega) \oplus_{\mathbf{L}_\varepsilon^2(\Omega)} \varepsilon^{-1} \text{curl} \mathbf{H}_{\Gamma_1}(\text{curl}, \Omega) \\ &= \nabla \mathbf{H}_{\Gamma_0}^1(\Omega) \oplus_{\mathbf{L}_\varepsilon^2(\Omega)} \mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega) \oplus_{\mathbf{L}_\varepsilon^2(\Omega)} \varepsilon^{-1} \text{curl} \mathbf{H}_{\Gamma_1}(\text{curl}, \Omega) \end{aligned}$$

hold (with continuous potential operators). Moreover, $\mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega)$ has finite dimension.

Combining Theorem 3.1 and Theorem 3.2 shows immediately the following.

Corollary 3.3 (regular Helmholtz decomposition with homogeneous mixed boundary conditions). *The $\mathbf{L}_\varepsilon^2(\Omega)$ -orthogonal regular Helmholtz decomposition*

$$\mathbf{L}_\varepsilon^2(\Omega) = \nabla \mathbf{H}_{\Gamma_0}^1(\Omega) \oplus_{\mathbf{L}_\varepsilon^2(\Omega)} \mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega) \oplus_{\mathbf{L}_\varepsilon^2(\Omega)} \varepsilon^{-1} \text{curl} \mathbf{H}_{\Gamma_1}^1(\Omega)$$

holds (with continuous potential operators) and $\mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega)$ has finite dimension. More precisely, any $E \in \mathbf{L}_\varepsilon^2(\Omega)$ may be $\mathbf{L}_\varepsilon^2(\Omega)$ -orthogonally (and regularly) decomposed into

$$E = \nabla u_\nabla + E_{\mathcal{H}} + \varepsilon^{-1} \text{curl} E_{\text{curl}}$$

with $u_\nabla \in \mathbf{H}_{\Gamma_0}^1(\Omega)$, $E_{\text{curl}} \in \mathbf{H}_{\Gamma_1}^1(\Omega)$, and $E_{\mathcal{H}} \in \mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega)$, and there exists a constant $c > 0$, independent of $E, u_\nabla, E_{\mathcal{H}}, E_{\text{curl}}$, such that

$$\begin{aligned} \|E_{\mathcal{H}}\|_{\mathbf{L}_\varepsilon^2(\Omega)} &\leq \|E\|_{\mathbf{L}_\varepsilon^2(\Omega)}, \\ c \|u_\nabla\|_{\mathbf{H}_{\Gamma_0}^1(\Omega)} &\leq \|\nabla u_\nabla\|_{\mathbf{L}_\varepsilon^2(\Omega)} \leq \|E\|_{\mathbf{L}_\varepsilon^2(\Omega)}, \\ c \|E_{\text{curl}}\|_{\mathbf{H}_{\Gamma_1}^1(\Omega)} &\leq \|\varepsilon^{-1} \text{curl} E_{\text{curl}}\|_{\mathbf{L}_\varepsilon^2(\Omega)} \leq \|E\|_{\mathbf{L}_\varepsilon^2(\Omega)}. \end{aligned}$$

4. COMPACT EMBEDDINGS

Our main result reads as follows:

Theorem 4.1 (compact embedding for vector fields with inhomogeneous mixed boundary conditions).

$$\widehat{\mathbf{H}}_{\Gamma_0}(\text{curl}, \Omega) \cap \varepsilon^{-1} \widehat{\mathbf{H}}_{\Gamma_1}(\text{div}, \Omega) \xrightarrow{\text{cpt}} \mathbf{L}^2(\Omega).$$

Proof. Let (E_ℓ) be a bounded sequence in $\widehat{\mathbf{H}}_{\Gamma_0}(\text{curl}, \Omega) \cap \varepsilon^{-1} \widehat{\mathbf{H}}_{\Gamma_1}(\text{div}, \Omega)$. By the Helmholtz decomposition in Corollary 3.3 we $\mathbf{L}_\varepsilon^2(\Omega)$ -orthogonally and regularly decompose

$$E_\ell = \nabla u_{\nabla,\ell} + E_{\mathcal{H},\ell} + \varepsilon^{-1} \text{curl} E_{\text{curl},\ell}$$

with $u_{\nabla,\ell} \in \mathbf{H}_{\Gamma_0}^1(\Omega)$, $E_{\text{curl},\ell} \in \mathbf{H}_{\Gamma_1}^1(\Omega)$, and $E_{\mathcal{H},\ell} \in \mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega)$, and there exists a constant $c > 0$ such that independent of E_{\dots} and for all ℓ

$$\|u_{\nabla,\ell}\|_{\mathbf{H}_{\Gamma_0}^1(\Omega)} + \|E_{\mathcal{H},\ell}\|_{\mathbf{L}_\varepsilon^2(\Omega)} + \|E_{\text{curl},\ell}\|_{\mathbf{H}_{\Gamma_1}^1(\Omega)} \leq c \|E_\ell\|_{\mathbf{L}_\varepsilon^2(\Omega)}.$$

As $\mathcal{H}_{\Gamma_0,\Gamma_1,\varepsilon}(\Omega)$ is finite dimensional we may assume (after extracting a subsequence) that $E_{\mathcal{H},\ell}$ converges strongly in $\mathbf{L}_\varepsilon^2(\Omega)$. Since $\mathbf{H}^1(\Omega) \xrightarrow{\text{cpt}} \mathbf{L}^2(\Omega)$ by Rellich's selection theorem, we may assume that also the regular potentials $u_{\nabla,\ell}$ and $E_{\text{curl},\ell}$ converge strongly in $\mathbf{L}^2(\Omega)$. Moreover, $u_{\nabla,\ell}|_\Gamma$ and $E_{\text{curl},\ell}|_\Gamma$ are bounded in $\mathbf{H}^{1/2}(\Gamma)$ by the (scalar) trace theorem, and thus we may assume by the compact embedding $\mathbf{H}^{1/2}(\Gamma) \xrightarrow{\text{cpt}} \mathbf{L}^2(\Gamma)$ that $u_{\nabla,\ell}|_\Gamma$ and $E_{\text{curl},\ell}|_\Gamma$ converge strongly in $\mathbf{L}^2(\Gamma)$. In particular, $u_{\nabla,\ell}|_{\Gamma_1}$ and $E_{\text{curl},\ell}|_{\Gamma_0}$ converge strongly in $\mathbf{L}^2(\Gamma_1)$ and $\mathbf{L}^2(\Gamma_0)$, respectively. After all this successively taking subsequences we obtain (using $\mathbf{L}_\varepsilon^2(\Omega)$ -orthogonality and the definition of the $\mathbf{L}^2(\Gamma_1)$ -traces of $\nu_{\Gamma_1} \varepsilon E_\ell$ and the $\mathbf{L}^2(\Gamma_0)$ -traces of $\tau_{\Gamma_0} E_\ell$ from Definition 2.1)

$$\begin{aligned} &\|\nabla(u_{\nabla,\ell} - u_{\nabla,k})\|_{\mathbf{L}_\varepsilon^2(\Omega)}^2 \\ &= \langle \nabla(u_{\nabla,\ell} - u_{\nabla,k}), E_\ell - E_k \rangle_{\mathbf{L}_\varepsilon^2(\Omega)} \\ &= -\langle u_{\nabla,\ell} - u_{\nabla,k}, \text{div} \varepsilon(E_\ell - E_k) \rangle_{\mathbf{L}^2(\Omega)} + \langle \sigma_{\Gamma_1}(u_{\nabla,\ell} - u_{\nabla,k}), \nu_{\Gamma_1} \varepsilon(E_\ell - E_k) \rangle_{\mathbf{L}^2(\Gamma_1)} \end{aligned}$$

$$\leq c\|u_{\nabla,\ell} - u_{\nabla,k}\|_{L^2(\Omega)} + c\|(u_{\nabla,\ell} - u_{\nabla,k})|_{\Gamma_1}\|_{L^2(\Gamma_1)} \rightarrow 0$$

and

$$\begin{aligned} & \|\varepsilon^{-1} \operatorname{curl}(E_{\operatorname{curl},\ell} - E_{\operatorname{curl},k})\|_{L^2_\varepsilon(\Omega)}^2 \\ &= \langle \varepsilon^{-1} \operatorname{curl}(E_{\operatorname{curl},\ell} - E_{\operatorname{curl},k}), E_\ell - E_k \rangle_{L^2_\varepsilon(\Omega)} \\ &= \langle E_{\operatorname{curl},\ell} - E_{\operatorname{curl},k}, \operatorname{curl}(E_\ell - E_k) \rangle_{L^2(\Omega)} + \langle \tau_{\Gamma_0}^\times(E_{\operatorname{curl},\ell} - E_{\operatorname{curl},k}), \tau_{\Gamma_0}(E_\ell - E_k) \rangle_{L^2(\Gamma_0)} \\ &\leq c\|E_{\operatorname{curl},\ell} - E_{\operatorname{curl},k}\|_{L^2(\Omega)} + c\|(E_{\operatorname{curl},\ell} - E_{\operatorname{curl},k})|_{\Gamma_0}\|_{L^2(\Gamma_0)} \rightarrow 0. \end{aligned}$$

Hence, (E_ℓ) contains a strongly $L^2_\varepsilon(\Omega)$ -convergent (and thus strongly $L^2(\Omega)$ -convergent) subsequence. \square

Remark 4.2 (compact embedding for vector fields with inhomogeneous mixed boundary conditions). *Theorem 4.1 even holds for weaker boundary data. For this, let $0 \leq s < 1/2$. Taking into account the compact embedding $H^{1/2}(\Gamma) \xrightarrow{\text{cpt}} H^s(\Gamma)$ and looking at the latter proof, we see that*

$$\{E \in H(\operatorname{curl}, \Omega) : \tau_{\Gamma_0} E \in H^{-s}(\Gamma_0)\} \cap \{E \in \varepsilon^{-1} H(\operatorname{div}, \Omega) : \nu_{\Gamma_1} \varepsilon E \in H^{-s}(\Gamma_1)\} \xrightarrow{\text{cpt}} L^2(\Omega).$$

5. APPLICATIONS

5.1. Friedrichs/Poincaré Type Estimates. A first application is the following estimate:

Theorem 5.1 (Friedrichs/Poincaré type estimate for vector fields with inhomogeneous mixed boundary conditions). *There exists a positive constant c such that for all vector fields E in $\widehat{H}_{\Gamma_0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1} \widehat{H}_{\Gamma_1}(\operatorname{div}, \Omega) \cap \mathcal{H}_{\Gamma_0, \Gamma_1, \varepsilon}(\Omega)^{\perp L^2_\varepsilon(\Omega)}$ it holds*

$$c\|E\|_{L^2_\varepsilon(\Omega)} \leq \|\operatorname{curl} E\|_{L^2(\Omega)} + \|\operatorname{div} \varepsilon E\|_{L^2(\Omega)} + \|\tau_{\Gamma_0} E\|_{L^2(\Gamma_0)} + \|\nu_{\Gamma_1} \varepsilon E\|_{L^2(\Gamma_1)}.$$

Proof. For a proof we use a standard compactness argument using Theorem 4.1. If the estimate was wrong, then there exists a sequence $(E_\ell) \in \widehat{H}_{\Gamma_0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1} \widehat{H}_{\Gamma_1}(\operatorname{div}, \Omega) \cap \mathcal{H}_{\Gamma_0, \Gamma_1, \varepsilon}(\Omega)^{\perp L^2_\varepsilon(\Omega)}$ with $\|E_\ell\|_{L^2_\varepsilon(\Omega)} = 1$ and

$$\|\operatorname{curl} E_\ell\|_{L^2(\Omega)} + \|\operatorname{div} \varepsilon E_\ell\|_{L^2(\Omega)} + \|\tau_{\Gamma_0} E_\ell\|_{L^2(\Gamma_0)} + \|\nu_{\Gamma_1} \varepsilon E_\ell\|_{L^2(\Gamma_1)} \rightarrow 0.$$

Thus, by Theorem 4.1 (after extracting a subsequence)

$$E_\ell \rightarrow E \quad \text{in} \quad \widehat{H}_{\Gamma_0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1} \widehat{H}_{\Gamma_1}(\operatorname{div}, \Omega) \cap \mathcal{H}_{\Gamma_0, \Gamma_1, \varepsilon}(\Omega)^{\perp L^2_\varepsilon(\Omega)} \quad (\text{strongly})$$

and $\operatorname{curl} E = 0$ and $\operatorname{div} \varepsilon E = 0$ (by testing). Moreover, for all $\Phi \in C_1^\infty(\Omega)$ and for all $\phi \in C_{\Gamma_0}^\infty(\Omega)$

$$\begin{aligned} \langle \operatorname{curl} \Phi, E_\ell \rangle_{L^2(\Omega)} - \langle \Phi, \operatorname{curl} E_\ell \rangle_{L^2(\Omega)} &= \langle \tau_{\Gamma_0}^\times \Phi, \tau_{\Gamma_0} E_\ell \rangle_{L^2(\Gamma_0)} \leq c\|\tau_{\Gamma_0} E_\ell\|_{L^2(\Gamma_0)} \rightarrow 0, \\ \langle \nabla \phi, \varepsilon E_\ell \rangle_{L^2(\Omega)} + \langle \phi, \operatorname{div} \varepsilon E_\ell \rangle_{L^2(\Omega)} &= \langle \sigma_{\Gamma_1} \phi, \nu_{\Gamma_1} \varepsilon E_\ell \rangle_{L^2(\Gamma_1)} \leq c\|\nu_{\Gamma_1} \varepsilon E_\ell\|_{L^2(\Gamma_1)} \rightarrow 0, \end{aligned}$$

cf. Definition 2.1, implying

$$\langle \operatorname{curl} \Phi, E \rangle_{L^2(\Omega)} = \langle \nabla \phi, \varepsilon E \rangle_{L^2(\Omega)} = 0.$$

Hence, $E \in H_{\Gamma_0, 0}(\operatorname{curl}, \Omega) \cap \varepsilon^{-1} H_{\Gamma_1, 0}(\operatorname{div}, \Omega) = \mathcal{H}_{\Gamma_0, \Gamma_1, \varepsilon}(\Omega)$ by [4, Theorem 4.7] (weak and strong homogeneous boundary conditions coincide). This shows $E = 0$ as $E \perp_{L^2_\varepsilon(\Omega)} \mathcal{H}_{\Gamma_0, \Gamma_1, \varepsilon}(\Omega)$, in contradiction to $1 = \|E_\ell\|_{L^2_\varepsilon(\Omega)} \rightarrow \|E\|_{L^2_\varepsilon(\Omega)} = 0$. \square

Remark 5.2 (Friedrichs/Poincaré type estimate for vector fields with inhomogeneous mixed boundary conditions). *As in Remark 4.2 there are corresponding generalised Friedrichs/Poincaré type estimates for weaker boundary data, where the $L^2(\Gamma_{0/1})$ -spaces and norms are replaced by $H^{-s}(\Gamma_{0/1})$ -spaces and norms.*

5.2. A div-curl Lemma. Another immediate consequence is a div-curl-lemma.

Theorem 5.3 (div-curl lemma for vector fields with inhomogeneous mixed boundary conditions).

Let (E_n) and (H_n) be bounded sequences in $\widehat{H}_{\Gamma_0}(\text{curl}, \Omega)$ and $\widehat{H}_{\Gamma_1}(\text{div}, \Omega)$, respectively. Then there exist $E \in \widehat{H}_{\Gamma_0}(\text{curl}, \Omega)$ and $H \in \widehat{H}_{\Gamma_1}(\text{div}, \Omega)$ as well as subsequences, again denoted by (E_n) and (H_n) , such that $E_n \rightharpoonup E$ in $\widehat{H}_{\Gamma_0}(\text{curl}, \Omega)$ and $H_n \rightharpoonup H$ in $\widehat{H}_{\Gamma_1}(\text{div}, \Omega)$ as well as

$$\langle E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle E, H \rangle_{L^2(\Omega)}.$$

Proof. We follow in closed lines the proof of [14, Theorem 3.1]. Let (E_n) and (H_n) be as stated. First, we pick subsequences, again denoted by (E_n) and (H_n) , and E and H , such that $E_n \rightharpoonup E$ in $\widehat{H}_{\Gamma_0}(\text{curl}, \Omega)$ and $H_n \rightharpoonup H$ in $\widehat{H}_{\Gamma_1}(\text{div}, \Omega)$. In particular,

$$(1) \quad \nu_{\Gamma_1} H_n \rightharpoonup \nu_{\Gamma_1} H \quad \text{in} \quad L^2(\Gamma_1).$$

To see (1), let $\nu_{\Gamma_1} H_n \rightharpoonup H_{\Gamma_1}$ in $L^2(\Gamma_1)$. Since for all $\phi \in H_{\Gamma_0}^1(\Omega)$

$$\begin{aligned} \langle \sigma_{\Gamma_1} \phi, H_{\Gamma_1} \rangle_{L^2(\Gamma_1)} &\leftarrow \langle \sigma_{\Gamma_1} \phi, \nu_{\Gamma_1} H_n \rangle_{L^2(\Gamma_1)} = \langle \nabla \phi, H_n \rangle_{L^2(\Omega)} + \langle \phi, \text{div} H_n \rangle_{L^2(\Omega)} \\ &\rightarrow \langle \nabla \phi, H \rangle_{L^2(\Omega)} + \langle \phi, \text{div} H \rangle_{L^2(\Omega)}, \end{aligned}$$

we get $H \in \widehat{H}_{\Gamma_1}(\text{div}, \Omega)$ and $\nu_{\Gamma_1} H = H_{\Gamma_1}$. Moreover, $\langle \sigma_{\Gamma_1} \phi, \nu_{\Gamma_1} H_n \rangle_{L^2(\Gamma_1)} \rightarrow \langle \sigma_{\Gamma_1} \phi, \nu_{\Gamma_1} H \rangle_{L^2(\Gamma_1)}$. As $\sigma_{\Gamma_1} H_{\Gamma_0}^1(\Omega)$ is dense in $L^2(\Gamma_1)$ and $(\langle \cdot, \nu_{\Gamma_1} H_n \rangle_{L^2(\Gamma_1)})$ is uniformly bounded with respect to n we obtain (1).

By Theorem 3.2 we have the orthogonal Helmholtz decomposition

$$\widehat{H}_{\Gamma_0}(\text{curl}, \Omega) \ni E_n = \nabla u_n + \widetilde{E}_n$$

with $u_n \in H_{\Gamma_0}^1(\Omega)$ and $\widetilde{E}_n \in \widehat{H}_{\Gamma_0}(\text{curl}, \Omega) \cap H_{\Gamma_1,0}(\text{div}, \Omega)$ as $\nabla H_{\Gamma_0}^1(\Omega) \subset H_{\Gamma_0,0}(\text{curl}, \Omega) \subset \widehat{H}_{\Gamma_0}(\text{curl}, \Omega)$. By orthogonality and the Friedrichs/Poincaré estimate, (u_n) is bounded in $H_{\Gamma_0}^1(\Omega)$ and hence contains a strongly $L^2(\Omega)$ -convergent subsequence, again denoted by (u_n) . (For $\Gamma_0 = \emptyset$ we may have to add a constant to each u_n .) Moreover, as $(u_n|_{\Gamma})$ is bounded in $H^{1/2}(\Gamma) \xrightarrow{\text{cpt}} L^2(\Gamma)$ we may assume that $(u_n|_{\Gamma})$ converges strongly in $L^2(\Gamma)$. In particular, $(\sigma_{\Gamma_1} u_n) = (u_n|_{\Gamma_1})$ converges strongly in $L^2(\Gamma_1)$. The sequence (\widetilde{E}_n) is bounded in $\widehat{H}_{\Gamma_0}(\text{curl}, \Omega) \cap H_{\Gamma_1,0}(\text{div}, \Omega)$ by orthogonality and since $\text{curl} \widetilde{E}_n = \text{curl} E_n$ and $\tau_{\Gamma_0} \widetilde{E}_n = \tau_{\Gamma_0} E_n$. Theorem 4.1 yields a strongly $L^2(\Omega)$ -convergent subsequence, again denoted by (\widetilde{E}_n) . Hence, there exist $u \in H_{\Gamma_0}^1(\Omega)$ and $\widetilde{E} \in \widehat{H}_{\Gamma_0}(\text{curl}, \Omega) \cap H_{\Gamma_1,0}(\text{div}, \Omega)$ such that $u_n \rightharpoonup u$ in $H_{\Gamma_0}^1(\Omega)$ and $u_n \rightarrow u$ in $L^2(\Omega)$ and $\sigma_{\Gamma_1} u_n \rightarrow \sigma_{\Gamma_1} u$ in $L^2(\Gamma_1)$ as well as $\widetilde{E}_n \rightharpoonup \widetilde{E}$ in $\widehat{H}_{\Gamma_0}(\text{curl}, \Omega) \cap H_{\Gamma_1,0}(\text{div}, \Omega)$ and $\widetilde{E}_n \rightarrow \widetilde{E}$ in $L^2(\Omega)$. Finally, we compute

$$\begin{aligned} \langle E_n, H_n \rangle_{L^2(\Omega)} &= \langle \nabla u_n, H_n \rangle_{L^2(\Omega)} + \langle \widetilde{E}_n, H_n \rangle_{L^2(\Omega)} \\ &= -\langle u_n, \text{div} H_n \rangle_{L^2(\Omega)} + \langle \sigma_{\Gamma_1} u_n, \nu_{\Gamma_1} H_n \rangle_{L^2(\Gamma_1)} + \langle \widetilde{E}_n, H_n \rangle_{L^2(\Omega)} \\ &\rightarrow -\langle u, \text{div} H \rangle_{L^2(\Omega)} + \langle \sigma_{\Gamma_1} u, \nu_{\Gamma_1} H \rangle_{L^2(\Gamma_1)} + \langle \widetilde{E}, H \rangle_{L^2(\Omega)} \\ &= \langle \nabla u, \text{div} H \rangle_{L^2(\Omega)} + \langle \widetilde{E}, H \rangle_{L^2(\Omega)} = \langle E, H \rangle_{L^2(\Omega)}, \end{aligned}$$

since indeed $E = \nabla u + \widetilde{E}$ holds by the weak convergence. \square

Remark 5.4 (div-curl lemma for vector fields with inhomogeneous mixed boundary conditions).

As in Remark 4.2 and Remark 5.2 there are corresponding generalised div-curl lemmas for weaker boundary data, where the $L^2(\Gamma_{0/1})$ -spaces and norms are replaced by $H^{-s}(\Gamma_{0/1})$ -spaces and norms.

5.3. Maxwell's Equations with Mixed Impedance Type Boundary Conditions. Let $\varepsilon,$

μ be admissible and time-independent matrix fields, and let $T, k \in \mathbb{R}_+$. In $I \times \Omega$ with $I := (0, T)$ we consider Maxwell's equations with mixed tangential and impedance boundary conditions

$$(2a) \quad \partial_t E - \varepsilon^{-1} \text{curl} H = F \in L^2(I, L_\varepsilon^2(\Omega)), \quad (\text{Ampère/Maxwell law})$$

$$(2b) \quad \partial_t H + \mu^{-1} \text{curl} E = G \in L^2(I, L_\mu^2(\Omega)), \quad (\text{Faraday/Maxwell law})$$

$$(2c) \quad \text{div} \varepsilon E = \rho, \quad (\text{Gauß law})$$

$$(2d) \quad \text{div} \mu H = 0, \quad (\text{Gauß law for magnetism})$$

$$\begin{aligned}
 (2e) \quad & \tau_{\Gamma_0} E = 0, & & \text{(perfect conductor bc)} \\
 (2f) \quad & \nu_{\Gamma_0} H = f, & & \text{(normal trace bc)} \\
 (2g) \quad & \tau_{\Gamma_1} E + k\tau_{\Gamma_1}^\times H = 0, & & \text{(impedance bc)} \\
 (2h) \quad & E(0) = E_0 \in \mathbb{L}_\varepsilon^2(\Omega), & & \text{(electric initial value)} \\
 (2i) \quad & H(0) = H_0 \in \mathbb{L}_\mu^2(\Omega). & & \text{(magnetic initial value)}
 \end{aligned}$$

Note that the impedance boundary condition, also called Leontovich boundary condition, is of Robin type and that the impedance is given by $\lambda = 1/k = \sqrt{\varepsilon/\mu}$, if $\varepsilon, \mu \in \mathbb{R}_+$.

Despite of other recent and very powerful approaches such as the concept of “evolutionary equations”, see the pioneering work of Rainer Picard, e.g., [20, 10], one can use classical semigroup theory for solving the Maxwell system (2).

We will split the system (2) into two static systems and a dynamic system. For simplicity we set $\varepsilon = \mu = 1$ and $F = G = 0$. The static systems are

$$\begin{aligned}
 (3a) \quad & \operatorname{curl} E = 0, & \operatorname{curl} H = 0, \\
 (3b) \quad & \operatorname{div} E = \rho, & \operatorname{div} H = 0, \\
 (3c) \quad & \tau_{\Gamma_0} E = 0, & \nu_{\Gamma_0} H = f, \\
 (3d) \quad & \tau_{\Gamma_1} E = -kg, & \tau_{\Gamma_1}^\times H = g,
 \end{aligned}$$

where g is any suitable tangential vector field in $\mathbb{L}^2(\Gamma_1)$. For simplicity we put $g = 0$, then these two systems are solvable by [2, Theorem 5.6]. However, the same result also gives conditions for which $g \neq 0$ this system is solvable. The dynamic system is

$$\begin{aligned}
 (4a) \quad & \partial_t E = \operatorname{curl} H, \\
 (4b) \quad & \partial_t H = -\operatorname{curl} E, \\
 (4c) \quad & \operatorname{div} E = 0, \\
 (4d) \quad & \operatorname{div} H = 0, \\
 (4e) \quad & \nu_{\Gamma_0} H = 0, \\
 (4f) \quad & \tau_{\Gamma_0} E = 0, \\
 (4g) \quad & \tau_{\Gamma_1} E + k\tau_{\Gamma_1}^\times H = 0.
 \end{aligned}$$

The initial conditions for the dynamic system are $E(0) = E_0 - E_{\text{stat}}$ and $H(0) = H_0 - H_{\text{stat}}$, where E_{stat} and H_{stat} are the solutions of the two static systems (3). We can write (4a) and (4b) as

$$\partial_t \begin{bmatrix} E \\ H \end{bmatrix} = \underbrace{\begin{bmatrix} 0 & \operatorname{curl} \\ -\operatorname{curl} & 0 \end{bmatrix}}_{=: A_0} \begin{bmatrix} E \\ H \end{bmatrix},$$

and the boundary conditions (4f) and (4g) shall be covered by the domain of A_0 :

$$\operatorname{dom} A_0 := \left\{ (E, H) \in \widehat{H}_\Gamma(\operatorname{curl}, \Omega) \times \widehat{H}_{\Gamma_1}(\operatorname{curl}, \Omega) \mid \tau_{\Gamma_0} E = 0, \tau_{\Gamma_1} E + k\tau_{\Gamma_1}^\times H = 0 \right\}.$$

Here, we did ignore the equations $\operatorname{div} E = 0$, $\operatorname{div} H = 0$ and $\nu_{\Gamma_0} H = 0$. However, A_0 is a generator of a C_0 -semigroup by [22, Example 8.10] or [25, Section 5], where the input function is $u = 0$. (In these sources they regard boundary control systems and system nodes, respectively. One condition of those concepts is that the system with $u = 0$ is described by a generator of a C_0 -semigroup). The next lemma provides a tool to respect the remaining conditions of (4) as well.

Lemma 5.5. *Let $T(\cdot)$ be a C_0 -semigroup on a Banach space X , and let A be its generator. Then every subspace $V \supseteq \operatorname{ran} A$ is invariant under $T(\cdot)$. Moreover, $A|_V$ generates the strongly continuous semigroup $T_V(\cdot) := T(\cdot)|_V$, if V is additionally closed in X .*

Proof. Let $t \geq 0$ and let $x \in V$. Then $\operatorname{ran} A \ni A \int_0^t T(s)x ds = T(t)x - x$ and hence $T(t)x \in V$. The remaining assertion follows from [6, Chapter II, Section 2.3]. \square

Therefore, it is left to show that the remaining conditions establish a closed and invariant subspace under the semigroup T_0 generated by A_0 or contains $\text{ran } A_0$. Note that by Theorem 3.1

$$\begin{aligned} S &:= \{(E, H) \mid \text{div } E = 0, \text{div } H = 0, \nu_{\Gamma_0} H = 0\} \\ &= \mathbf{H}_0(\text{div}, \Omega) \times \mathbf{H}_{\Gamma_0, 0}(\text{div}, \Omega) \\ &= (\text{curl } \mathbf{H}(\text{curl}, \Omega) \times \text{curl } \mathbf{H}_{\Gamma_0}(\text{curl}, \Omega)) \oplus (\mathcal{H}_{\Gamma, \emptyset}(\Omega) \times \mathcal{H}_{\Gamma_1, \Gamma_0}(\Omega)). \end{aligned}$$

This space is closed as the intersection of kernels of closed operators. Clearly, $\mathcal{H}_{\Gamma, \emptyset}(\Omega) \times \mathcal{H}_{\Gamma_1, \Gamma_0}(\Omega)$ is invariant under T_0 , since every $(E, H) \in \mathcal{H}_{\Gamma, \emptyset}(\Omega) \times \mathcal{H}_{\Gamma_1, \Gamma_0}(\Omega)$ is a constant in time solution of the system (4), i.e.,

$$T_0(t) \begin{bmatrix} E \\ H \end{bmatrix} = \begin{bmatrix} E \\ H \end{bmatrix}.$$

By

$$\text{curl } \mathbf{H}(\text{curl}, \Omega) \times \text{curl } \mathbf{H}_{\Gamma_0}(\text{curl}, \Omega) = \begin{bmatrix} 0 & \text{curl} \\ -\text{curl} & 0 \end{bmatrix} (\mathbf{H}_{\Gamma_0}(\text{curl}, \Omega) \times \mathbf{H}(\text{curl}, \Omega)) \supseteq \text{ran } A_0$$

and Lemma 5.5 we have that also $\text{curl } \mathbf{H}(\text{curl}, \Omega) \times \text{curl } \mathbf{H}_{\Gamma_0}(\text{curl}, \Omega)$ is invariant under T_0 . Hence, Lemma 5.5 and Theorem 4.1 imply the next theorem.

Theorem 5.6. $A := A_0|_S$ is a generator of a C_0 -semigroup and

$$\text{dom } A \subseteq (\widehat{\mathbf{H}}_{\Gamma}(\text{curl}, \Omega) \cap \mathbf{H}(\text{div}, \Omega)) \times (\widehat{\mathbf{H}}_{\Gamma_1}(\text{curl}, \Omega) \cap \mathbf{H}_{\Gamma_0}(\text{div}, \Omega)) \xrightarrow{\text{cpt}} \mathbf{L}^2(\Omega).$$

Consequently, every resolvent operator of A is compact.

If $\mathcal{H}_{\Gamma, \emptyset}(\Omega) = \{0\}$ and $\mathcal{H}_{\Gamma_1, \Gamma_0}(\Omega) = \{0\}$, then 0 is in the resolvent set of A and A^{-1} is compact. Alternatively, we can further restrict A to $\mathcal{H}_{\Gamma, \emptyset}(\Omega)^{\perp \mathbf{L}^2(\Omega)} \times \mathcal{H}_{\Gamma_1, \Gamma_0}(\Omega)^{\perp \mathbf{L}^2(\Omega)}$. This would also match our separation of static solutions and dynamic solutions, since solutions with initial condition in $\mathcal{H}_{\Gamma, \emptyset}(\Omega) \times \mathcal{H}_{\Gamma_1, \Gamma_0}(\Omega)$ are constant in time.

5.4. Wave Equation with Mixed Impedance Type Boundary Conditions. For the scalar wave equation the situation is even simpler since traces of $\mathbf{H}^1(\Omega)$ -functions already belong to $\mathbf{L}^2(\Gamma)$, even to $\mathbf{H}^{1/2}(\Gamma)$. In $I \times \Omega$ we consider the wave equation in first order form (linear acoustics) with mixed scalar and impedance boundary conditions

$$\begin{aligned} \partial_t w - \text{div } v &= 0, \\ \partial_t v - \nabla w &= 0, \\ \sigma_{\Gamma_0} w &= 0, \\ \sigma_{\Gamma_1} w + k\nu_{\Gamma_1} v &= 0, \\ w(0) &= w_0 \in \mathbf{L}^2(\Omega), \\ v(0) &= v_0 \in \mathbf{L}^2(\Omega). \end{aligned}$$

We write the system as

$$\partial_t \begin{bmatrix} w \\ v \end{bmatrix} = \underbrace{\begin{bmatrix} 0 & \text{div} \\ \nabla & 0 \end{bmatrix}}_{=: A_0} \begin{bmatrix} w \\ v \end{bmatrix}$$

with

$$\text{dom } A_0 := \{(w, v) \in \mathbf{H}_{\Gamma_0}^1(\Omega) \times \widehat{\mathbf{H}}_{\Gamma_1}(\text{div}, \Omega) \mid \sigma_{\Gamma_1} w + k\nu_{\Gamma_1} v = 0\}.$$

As before, by [8, Theorem 4.4] or [22, Example 8.9], A_0 is a generator of C_0 -semigroup. Again, we want to separate the static solutions from the dynamic system. The static solutions are given by $\ker A_0$, which can be characterise by

$$\ker A_0 = \{0\} \times \mathbf{H}_{\Gamma_1, 0}(\text{div}, \Omega),$$

where we assumed $\Gamma_0 \neq \emptyset$, otherwise the first component can also be constant and the second component would be in $\widehat{\mathbf{H}}_{\Gamma_1, 0}(\text{div}, \Omega)$. By Theorem 3.2, the orthogonal complement of $\ker A_0$ is

$$S := \mathbf{L}^2(\Omega) \times \nabla \mathbf{H}_{\Gamma_0}^1(\Omega).$$

Note that S contains $\text{ran } A_0$ and is therefore (by Lemma 5.5) an invariant subspace under the semigroup generated by A_0 . Moreover, note that $\nabla H_{\Gamma_0}^1(\Omega) \subseteq H_{\Gamma_0,0}(\text{curl}, \Omega)$ and that S is closed. Hence, Lemma 5.5 and Theorem 4.1 imply the next theorem.

Theorem 5.7. $A := A_0|_S$ is a generator of C_0 -semigroup and

$$\text{dom } A \subseteq H_{\Gamma_0}^1(\Omega) \times (\widehat{H}_{\Gamma_1}(\text{div}, \Omega) \cap H_{\Gamma_0,0}(\text{curl}, \Omega)) \xrightarrow{\text{cpt}} L^2(\Omega).$$

Consequently, every resolvent operator of A is compact.

Alternatively, we can also regard the classical formulation of the wave equation and see that it is necessary for the second component in our formulation to be in $\nabla H_{\Gamma_0}^1(\Omega)$, if we want the solutions to correspond.

REFERENCES

- [1] D.N. Arnold and K. Hu. Complexes from complexes. *arXiv*, <https://arxiv.org/abs/2005.12437>, 2020.
- [2] S. Bauer, D. Pauly, and M. Schomburg. The Maxwell compactness property in bounded weak Lipschitz domains with mixed boundary conditions. *SIAM J. Math. Anal.*, 48(4):2912–2943, 2016.
- [3] S. Bauer, D. Pauly, and M. Schomburg. Weck’s selection theorem: The Maxwell compactness property for bounded weak Lipschitz domains with mixed boundary conditions in arbitrary dimensions. *arXiv*, <https://arxiv.org/abs/1809.01192>, 2018.
- [4] S. Bauer, D. Pauly, and M. Schomburg. Weck’s selection theorem: The Maxwell compactness property for bounded weak Lipschitz domains with mixed boundary conditions in arbitrary dimensions. *Maxwell’s Equations: Analysis and Numerics (Radon Series on Computational and Applied Mathematics 24)*, De Gruyter, pages 77–104, 2019.
- [5] M. Costabel. A remark on the regularity of solutions of Maxwell’s equations on Lipschitz domains. *Math. Methods Appl. Sci.*, 12(4):365–368, 1990.
- [6] K. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt.
- [7] F. Jochmann. A compactness result for vector fields with divergence and curl in $L^q(\Omega)$ involving mixed boundary conditions. *Appl. Anal.*, 66:189–203, 1997.
- [8] Mikael Kurula and Hans Zwart. Linear wave systems on n -D spatial domains. *Internat. J. Control*, 88(5):1063–1077, 2015.
- [9] R. Leis. *Initial Boundary Value Problems in Mathematical Physics*. Teubner, Stuttgart, 1986.
- [10] D. McGhee, R. Picard, S. Trostorff, and M. Waurick. *A Primer for a Secret Shortcut to PDEs of Mathematical Physics*. Birkhäuser, 2020.
- [11] F. Osterbrink and D. Pauly. Time-harmonic electro-magnetic scattering in exterior weak lipschitz domains with mixed boundary conditions. *Maxwell’s Equations: Analysis and Numerics (Radon Series on Computational and Applied Mathematics 24)*, De Gruyter, pages 341–382, 2019.
- [12] F. Osterbrink and D. Pauly. Low frequency asymptotics and electro-magneto-statics for time-harmonic Maxwell’s equations in exterior weak Lipschitz domains with mixed boundary conditions. *SIAM J. Math. Anal.*, 52(5):4971–5000, 2020.
- [13] D. Pauly. On the Maxwell constants in 3D. *Math. Methods Appl. Sci.*, 40(2):435–447, 2017.
- [14] D. Pauly. A global div-curl-lemma for mixed boundary conditions in weak Lipschitz domains and a corresponding generalized A_0^* - A_1 -lemma in Hilbert spaces. *Analysis (Munich)*, 39(2):33–58, 2019.
- [15] D. Pauly. On the Maxwell and Friedrichs/Poincaré constants in ND. *Math. Z.*, 293(3-4):957–987, 2019.
- [16] D. Pauly. Solution theory, variational formulations, and functional a posteriori error estimates for general first order systems with applications to electro-magneto-statics and more. *Numer. Funct. Anal. Optim.*, 41(1):16–112, 2020.
- [17] D. Pauly and W. Zulehner. The divDiv-complex and applications to biharmonic equations. *Appl. Anal.*, 99(9):1579–1630, 2020.
- [18] D. Pauly and W. Zulehner. The elasticity complex: Compact embeddings and regular decompositions. *arXiv*, <https://arxiv.org/abs/2001.11007>, 2020.
- [19] R. Picard. An elementary proof for a compact imbedding result in generalized electromagnetic theory. *Math. Z.*, 187:151–164, 1984.
- [20] R. Picard. A structural observation for linear material laws in classical mathematical physics. *Math. Methods Appl. Sci.*, 32(14):1768–1803, 2009.
- [21] R. Picard, N. Weck, and K.-J. Witsch. Time-harmonic Maxwell equations in the exterior of perfectly conducting, irregular obstacles. *Analysis (Munich)*, 21:231–263, 2001.
- [22] N. Skrepek. Well-posedness of linear first order port-hamiltonian systems on multidimensional spatial domains. *Evolution Equations & Control Theory*, (2163-2480.2019.0.91), 2020.
- [23] C. Weber. A local compactness theorem for Maxwell’s equations. *Math. Methods Appl. Sci.*, 2:12–25, 1980.

- [24] N. Weck. Maxwell's boundary value problems on Riemannian manifolds with nonsmooth boundaries. *J. Math. Anal. Appl.*, 46:410–437, 1974.
 - [25] G. Weiss and O. J. Staffans. Maxwell's equations as a scattering passive linear system. *SIAM J. Control Optim.*, 51(5):3722–3756, 2013.
 - [26] K.-J. Witsch. A remark on a compactness result in electromagnetic theory. *Math. Methods Appl. Sci.*, 16:123–129, 1993.
-

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, GERMANY
Email address, Dirk Pauly: **dirk.pauly@uni-due.de**

FAKULTÄT FÜR MATHEMATIK UND NATURWISSENSCHAFTEN, BERGISCHE UNIVERSITÄT WUPPERTAL, GERMANY
Email address, Nathanael Skrepek: **skrepek@uni-wuppertal.de**