

# GENERALIZATION OF A CRITERION FOR SEMISTABLE VECTOR BUNDLES

INDRANIL BISWAS AND GEORG HEIN

ABSTRACT. It is known that a vector bundle  $E$  on a smooth projective curve  $Y$  defined over an algebraically closed field is semistable if and only if there is a vector bundle  $F$  on  $Y$  such that both  $H^0(X, E \otimes F)$  and  $H^1(X, E \otimes F)$  vanishes. We extend this criterion for semistability to vector bundles on curves defined over perfect fields. Let  $X$  be a geometrically irreducible smooth projective curve defined over a perfect field  $k$ , and let  $E$  be a vector bundle on  $X$ . We prove that  $E$  is semistable if and only if there is a vector bundle  $F$  on  $X$  such that  $H^i(X, E \otimes F) = 0$  for all  $i$ . We also give an explicit bound for the rank of  $F$ .

## 1. INTRODUCTION

A theorem due to Faltings says that a vector bundle  $E$  on a smooth projective curve  $Y$  defined over an algebraically closed field of characteristic zero is semistable if and only if there is a vector bundle  $F$  on  $Y$  such that both  $H^i(X, E \otimes F) = 0$  for all  $i$  [3, p. 514, Theorem 1.2]. It is known that this criterion for semistability extends to vector bundles on smooth projective curves defined over an algebraically closed fields of positive characteristic. (See [5], [1] for related results.)

Our aim here is to investigate this criterion for curves defined over finite fields and more generally over perfect fields. We prove the following theorem.

**Theorem 1.1.** *Let  $X$  be a geometrically irreducible smooth projective curve defined over a perfect field  $k$ . A vector bundle  $E$  over  $X$  is semistable if and only if there is a vector bundle  $F$  over  $X$  such that  $H^i(X, E \otimes F) = 0$  for all  $i$ .*

We also produce an effective bound for the rank of  $F$  in Theorem 1.1. More precisely, given nonnegative integers  $g$  and  $r$ , and an integer  $d$ , there is an explicit integer  $R(g, r, d)$  such that for any triple  $(k, X, E)$ , where

- $k$  is a perfect field,
- $X$  is a geometrically irreducible smooth projective curve of genus  $g$  defined over  $k$ , and
- $E$  is a vector bundle over  $X$  of rank  $r$  and degree  $d$ ,

the vector bundle  $E$  is semistable if and only if there is a vector bundle  $F$  over  $X$  of rank  $R(g, r, d)$  such that  $H^i(X, E \otimes F) = 0$  for all  $i$ . (See Theorem 3.1.)

---

*Date:* April 25, 2008.

Let  $X$  be a scheme defined over a field  $k$ , and let  $D \subset X$  be an effective divisor. Then there is a geometric point of  $X$  that lies outside  $D$ . Corollary 2.5 bounds the degree of the field extension  $K/k$  such that  $D(K) \subsetneq X(K)$ . This is used in the proof of Theorem 1.1.

## 2. RATIONAL POINTS OUTSIDE A GIVEN HYPERSURFACE

Let  $k$  be any field. The algebraic closure of  $k$  will be denoted by  $\bar{k}$ .

**Lemma 2.1.** *Let  $D \subset \mathbb{A}_k^n$  be an effective divisor defined over  $\bar{k}$ . Given any field extension  $K/k$  such  $K$  has more than  $\deg(D)$  elements, there exists a  $K$ -rational point in  $\mathbb{A}_K^n$  that lies outside  $D$ .*

*Proof.* One follows the proof of Proposition 1.3(a) in [6, p. 4] almost word for word simply replacing “infinite field” by “field with more than  $\deg(D)$  elements”: We assume that  $D$  is given by the polynomial  $F \in \bar{k}[X_1, \dots, X_n]$ , and proceed by induction over  $n$ . For  $n = 1$  it is the statement that a polynomial of degree  $d$  cannot have more than  $d$  zeros. Now assume that  $X_n$  occurs in  $F$  and write

$$F = \varphi_0 + \varphi_1 X_n + \dots + \varphi_t X_n^t,$$

where  $\varphi_i \in \bar{k}[X_1, \dots, X_{n-1}]$  and  $\varphi_t \neq 0$ . Since  $t$  and  $\deg(\varphi_t)$  are both at most  $\deg(D)$ , we conclude from the induction hypothesis the existence of a point  $(x_1, \dots, x_{n-1}) \in K^{n-1}$  such that  $\varphi_t(x_1, \dots, x_{n-1}) \neq 0$ . Now the polynomial  $X_n \mapsto F(x_1, \dots, x_{n-1}, X_n)$  has at most  $t$  zeros.  $\square$

**Lemma 2.2.** *Let  $D \subset \mathbb{P}_k^n$  be an effective divisor in projective space  $\mathbb{P}^n$  defined over  $\bar{k}$ . Then for any extension  $K/k$  such that  $K$  has at least  $\deg(D)$  elements, there is a  $K$ -rational point in  $\mathbb{P}^n(K)$  that lies outside  $D(K)$ .*

*Proof.* Again we proceed by induction on  $n$ . The case  $n = 1$  is obvious. For  $n > 1$ , we consider the pencil of hyperplanes defined over  $K$  passing through a codimension two linear subspace. Since there are more than  $\deg(D)$  of these hyperplanes, the union of all these hyperplanes cannot be contained in  $D$ . Thus, there exists a hyperplane  $H \cong \mathbb{P}^{n-1} \subset \mathbb{P}^n$  that intersects  $D$  properly. Now the proof is completed by the induction hypothesis.  $\square$

Let  $\text{Grass}(m, n)$  be the Grassmannian of  $m$ -dimensional linear subspaces of  $k^n$ . Let

$$(1) \quad \iota : \text{Grass}(m, n) \longrightarrow \mathbb{P} := \mathbb{P}^{\binom{n}{m}-1}$$

be the Plücker embedding. By an hypersurface of degree  $d$  on  $\text{Grass}(m, n)$  we will mean one from the complete linear system  $|\iota^* \mathcal{O}_{\mathbb{P}}(d)|$ .

**Lemma 2.3.** *Let  $D \subset \text{Grass}(m, n)$  be a hypersurface in the Grassmannian. If a field extension  $K/k$  has more than  $m \cdot \deg(D)$  elements, then there is a  $K$ -rational point of  $\text{Grass}(m, n)(K)$  that is not contained in  $D(K)$ .*

*Proof.* We consider the dense open cell in the Grassmannian given by the open immersion  $j : \mathbb{A}^{m \cdot (n-m)} \longrightarrow \text{Grass}(m, n)$  defined by

$$(a_{i,j})_{i=1,\dots,m \ j:=m+1,\dots,n} \longmapsto \text{span} \begin{pmatrix} 1 & 0 & \cdots & 0 & a_{1,m+1} & a_{1,m+2} & \cdots & a_{1,n} \\ 0 & 1 & \cdots & 0 & a_{2,m+1} & a_{2,m+2} & \cdots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & a_{m,m+1} & a_{m,m+2} & \cdots & a_{m,n} \end{pmatrix}$$

The Plücker embedding (see (1)) restricted to  $\mathbb{A}^{m \cdot (n-m)}$  is given by the  $m \times m$ -minors of degree at most  $m$  of the above matrix. Therefore,  $j^*D$  is a divisor of degree  $m \cdot \deg(D)$ . Now the proof is completed using Lemma 2.1.  $\square$

**Proposition 2.4.** *Let  $\mathcal{O}_X(H)$  be a globally generated ample line bundle on a projective scheme  $X$  of dimension  $n$  defined over  $k$ . Let  $D \subset X$  be an effective divisor  $D \subset X$ . Let  $K_1/k$  be a field extension that has more than  $\max\{(n+1)H^n, D.H^{n-1} - 1\}$  elements. Then there exists a field extension  $K_2/K_1$  with  $[K_2 : K_1] \leq H^n$ , such that there is a  $K_2$ -rational point of  $X(K_2)$  that does not lie in  $D(K_2)$ .*

*Proof.* We consider the short exact sequence of vector bundles

$$0 \longrightarrow W \longrightarrow H^0(L) \otimes \mathcal{O}_X \longrightarrow L \longrightarrow 0$$

over  $X$ . Let  $\text{Grass}_X(n+1, W)$  be the Grassmann bundle over  $X$  parameterizing all  $(n+1)$ -dimensional subspaces in the fibers of  $W$ . We have

$$\begin{aligned} \dim \text{Grass}_X(n+1, W) &= \dim(X) + (h^0(L) - n - 2)(n+1) \\ &= \dim(\text{Grass}(n+1, H^0(L))) - 1. \end{aligned}$$

We will show that the degree of the hypersurface  $\text{Grass}_X(n+1, W)$  in  $\text{Grass}(n+1, H^0(L))$  is  $H^n$ . To prove this, take any subspace  $U \subset H^0(X, L)$  of dimension  $n+2$ . The  $(n+1)$ -dimensional subspaces of  $U$  form a projective line  $\mathbb{P}_k^1$  in  $\text{Grass}(n+1, H^0(L))$ . The degree of the restriction of  $\iota^*\mathcal{O}_{\mathbb{P}^1}(1)$  (see (1)) to this  $\mathbb{P}_k^1$  is one. To compute the intersection number of the line with  $\text{Grass}_X(n+1, W)$  we may assume that  $H^0(X, L) = U$ . So it suffices to count the intersection of a line in  $\mathbb{P}(U)$  with the divisor  $X \subset \mathbb{P}(U)$ . Thus, we conclude that the hypersurface  $\text{Grass}_X(n+1, W) \subset \text{Grass}(n+1, H^0(L))$  is of degree  $H^n$ .

Now, using Lemma 2.3 and the assumption on  $K_1$  we conclude that there exists a  $K$ -point in  $\text{Grass}(n+1, H^0(X, L))$  not lying in  $\text{Grass}_X(n+1, W)$ . This yields a finite morphism  $X \xrightarrow{\pi} \mathbb{P}^n$  defined over  $K_1$ . Now  $\pi_*(D)$  is a divisor of degree  $D.H^{-1}$  on  $\mathbb{P}^n$ . Our assumption on the number of elements in  $K_1$  and Lemma 2.2 together imply that there is a  $K_1$ -rational point  $P$  in the complement of  $\pi_*(D)$  in  $\mathbb{P}^n$ . The morphism  $X_P \longrightarrow \text{Spec}(K_1)$  is finite of degree  $H^n$ , and it is defined over  $K_1$ . Thus, we find at least one point in  $X_P$  defined over a field  $K_2$  as in the statement of the proposition. This completes the proof of the proposition.  $\square$

Proposition 2.4 has the following corollary.

**Corollary 2.5.** *Given positive integers  $n, \alpha$  and  $\beta$ , define*

$$M(n, \alpha, \beta) := \alpha \lceil \log_2(\max\{(n+1)\alpha + 1, \beta\}) \rceil.$$

For any quadruple  $(k, X, H, D)$ , where

- $k$  is a field,
- $X$  is a projective scheme of dimension  $n$  defined over  $k$ ,
- $H \subset X$  is a base point-free ample hypersurface with  $H^n = \alpha$ , and
- $D \subset H$  is an effective divisor with  $D.H^{n-1} = \beta$ ,

there is a field extension  $K/k$  of degree  $[K : k] \leq M(n, \alpha, \beta)$  with the property that  $X(K)$  has a  $K$ -rational point that does not lie in  $D(K)$ .

If we restrict ourselves only to infinite fields, then  $M(n, \alpha, \beta)$  in Corollary 2.5 can be taken to be  $\alpha$ . If we fix a prime  $p$  and restrict ourselves only to fields of characteristic  $p$ , then  $M(n, \alpha, \beta)$  in Corollary 2.5 can be taken to be  $\alpha \lceil \log_p(\max\{(n+1)\alpha + 1, \beta\}) \rceil$ .

### 3. SEMISTABILITY CRITERION OVER PERFECT FIELDS

**Theorem 3.1.** *Let  $X$  be a geometrically irreducible smooth projective curve of genus  $g$  defined over a perfect field  $k$ . Fix a positive integer  $r$  and an integer  $d$ . Then there is an explicit positive integer  $R$  that depends only on  $r, d$  and  $g$  (in particular,  $R$  is independent of  $k$ ) with the following property: A vector bundle  $E$  over  $X$  of rank  $r$  and degree  $d$  is semistable if and only if there is a vector bundle  $F$  over  $X$  of rank  $R$  such that  $H^i(X, E \otimes F) = 0$  for all  $i$ .*

*Proof.* If  $E$  is not semistable, then clearly there is no  $F$  such that  $H^i(X, E \otimes F) = 0$  for all  $i$ . Let  $E$  be a semistable vector bundle over  $X$  of rank  $r$  and degree  $d$ . We will construct  $R$  and  $F$ .

The moduli space of semistable vector bundles over  $X$  of rank  $r'$  and degree  $d'$  will be denoted by  $\mathcal{U}_X(r', d')$ .

Let  $h := \gcd(r, d)$ . Furthermore, we set  $\bar{r} := \frac{r}{h}$ , and  $\bar{d} := \frac{d}{h}$ . For any integer  $n \geq 1$ , consider the morphism  $\mathcal{U}_X(n\bar{r}, n(\bar{r}(g-1) - \bar{d})) \rightarrow \mathcal{U}_X(nr\bar{r}, nr\bar{r}(g-1))$  defined by  $V \mapsto V \otimes E$ . Let  $\Theta_E$  denote the pull back of the natural theta divisor in  $\mathcal{U}_X(nr\bar{r}, nr\bar{r}(g-1))$  by this morphism. Therefore,  $\Theta_E \times_k \bar{k}$  consists of all semistable vector bundles  $W$  over  $X_{\bar{k}} = X \times_k \bar{k}$  of rank  $n\bar{r}$  and degree  $n(\bar{r}(g-1) - \bar{d})$  such that  $H^0(X_{\bar{k}}, W \otimes (E \otimes_k \bar{k})) \neq 0$ . (We note that a vector bundle  $V'$  over  $X$  is semistable if and only if the vector bundle  $V' \otimes_k \bar{k}$  over  $X_{\bar{k}}$  is semistable; see [4, p. 222].) The subscheme  $\Theta_E$  defined above is either an effective Cartier divisor in the complete linear system  $|h \cdot \Theta|$  or it is the entire moduli space  $\mathcal{U}_X(n\bar{r}, n(\bar{r}(g-1) - \bar{d}))$  (cf. [2, § 0.2.1]).

Popa showed that this is indeed a divisor in the linear system  $|h \cdot \Theta|$  for all  $n \geq \frac{r^2+1}{4}$  (see [7, p. 490, Theorem 5.3]). Let  $n$  be the smallest integer such that  $n \geq \frac{r^2+1}{4}$ . Consider the effective divisor  $\Theta_E \subset \mathcal{U}_X(n\bar{r}, n(\bar{r}(g-1) - \bar{d}))$  for  $n := \lceil \frac{r^2+1}{4} \rceil$ . By Corollary 2.5, there exists an integer  $M$  and a field extension  $K/k$  such that  $\Theta_E$  does not contain all  $K$ -rational points of  $\mathcal{U}_X(n\bar{r}, n(\bar{r}(g-1) - \bar{d}))$ .

The integer  $R$  in the statement of the theorem will be  $n\bar{r}M!$ .

Since  $\Theta_E$  does not contain all  $K$ -rational points of  $\mathcal{U}_X(n\bar{r}, n(\bar{r}(g-1) - \bar{d}))$ , there exists a vector bundle  $F_1$  of rank  $n\bar{r}$  defined over  $X_K = X \times_k K$ , where  $K/k$  is some Galois extension of degree dividing  $M!$ , such that

$$(2) \quad H^0(X_K, (E \otimes_k K) \otimes F_1) = 0 = H^1(X_K, (E \otimes_k K) \otimes F_1).$$

From (2) it follows that

$$(3) \quad H^0(X_K, (E \otimes_k K) \otimes \sigma^* F_1) = 0 = H^1(X_K, (E \otimes_k K) \otimes \sigma^* F_1)$$

for all  $\sigma \in \text{Gal}(K/k)$ .

Now we consider the direct sum

$$F_2 := \bigoplus_{\sigma \in \text{Gal}(K/k)} \sigma^* F_1.$$

This  $F_2$  is a vector bundle defined over  $X$ . From (3) it follows immediately that  $H^i(X, E \otimes F_2) = 0$  for all  $i$ . Also, the rank of  $F_2$  clearly divides  $n\bar{r}M!$ . Finally we set  $m := \frac{n\bar{r}M!}{\text{rk}(F_2)}$ , and  $F := F_2^{\oplus m}$ , and obtain the asserted vector bundle of rank  $R = n\bar{r}M!$ .  $\square$

#### REFERENCES

- [1] I. Biswas, G. Hein, *Parabolic Raynaud bundles*, Manuscr. Math. (in press).
- [2] J.-M. Drezet, M. S. Narasimhan, *Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques*, Invent. Math. **97** (1989) 53–94.
- [3] G. Faltings, *Stable  $G$ -bundles and projective connections*, Jour. Alg. Geom. **2** (1993) 507–568.
- [4] G. Harder, M. S. Narasimhan, *On the cohomology groups of moduli spaces of vector bundles on curves*, Math. Ann. **212** (1975) 215–248.
- [5] G. Hein, *Raynaud vector bundles*, preprint, math.AG/0706.3970.
- [6] E. Kunz, *Introduction to Commutative Algebra and Algebraic Geometry*, Birkhäuser, Boston, 1985.
- [7] M. Popa, *Dimension estimates for Hilbert schemes and effective base point freeness on moduli spaces of vector bundles on curves*, Duke Math Jour. **107** (2001) 469–495.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

*E-mail address:* indranil@math.tifr.res.in

UNIVERSITÄT DUISBURG-ESSEN, FACHBEREICH MATHEMATIK, 45117 ESSEN, GERMANY

*E-mail address:* georg.hein@uni-due.de