

KÜNNETH PROJECTORS FOR OPEN VARIETIES

SPENCER BLOCH AND HÉLÈNE ESNAULT

To Jacob Murre

ABSTRACT. We consider correspondences on smooth quasiprojective varieties U . An algebraic cycle inducing the Künneth projector onto $H^1(U)$ is constructed. Assuming normal crossings at infinity, the existence of relative motivic cohomology is shown to imply the independence of ℓ for traces of open correspondences.

1. INTRODUCTION

Let X be a smooth, projective algebraic variety over an algebraically closed field k , and let $H^*(X)$ denote a Weil cohomology theory. The existence of algebraic cycles on $X \times X$ inducing as correspondences the various Künneth projectors $\pi^i : H^*(X) \rightarrow H^i(X)$ is one of the standard conjectures of Grothendieck, [9], [10]. It is known in general only for the cases $i = 0, 1, 2d - 1, 2d$ where $d = \dim X$. The purpose of this note is to consider correspondences on smooth quasi-projective varieties U . In the first section we prove the existence of an “algebraic” Künneth projector $\pi^1 : H^*(U) \rightarrow H^1(U)$ assuming that U admits a smooth, projective completion X . The word algebraic is placed in quotes here because in fact the algebraic cycle on $X \times U$ inducing π^1 is not, as one might imagine, trivialized on $(X - U) \times U$. It is only partially trivialized. This partial trivialization is sufficient to define a class in $H_c^{2d-1}(U) \otimes H^1(U)$ giving the desired projection. Of course, our cycle on $X \times U$ will be trivialized on $(X \setminus U) \times V$ for $V \subset U$ suitably small nonempty open, but our method does not in any obvious way yield a full trivialization on $(X \setminus U) \times U$. We finish this first section with some comments on π^i for $i > 1$ and some speculation, mostly coming from discussions with A. Beilinson, on how these ideas might be applied to study the Milnor conjecture that the Galois cohomology ring of the function field $H^*(k(X), \mathbb{Z}/n\mathbb{Z})$ is generated by H^1 .

In the last section, we use the existence of relative motivic cohomology [11] to prove an integrality and independence of ℓ result for the

Date: Feb. 20, 2005.

trace of an algebraic correspondence Γ on $U \times U$. We are indebted to G. Laumon for pointing out that one may endeavor to prove this using results already in the literature ([3], [13], [14], and [6]) by reduction mod p and composition with a high power of Frobenius. Our objective in what follows is to show how techniques in motivic cohomology can apply to such questions, at least when the divisor at infinity has normal crossings.

When the Zariski closure of the correspondence stabilizes the various strata D_I at infinity (e.g. when the correspondence is the graph of Frobenius) then the trace on $H^*(U)$ is realized as an alternating sum of traces on $H^*(D_I)$. When in addition all the intersections with the diagonals are transverse, the contribution to the alternating sum coming from points lying off U cancels, and the trace on $H^*(U)$ is just the sum of the fixed points on U .

We would like to acknowledge helpful correspondence with A. Beilinson, M. Levine, J. Murre, T. Saito, and V. Srinivas. We thank G. Laumon and L. Lafforgue for explaining to us [6].

2. THE FIRST KÜNNETH COMPONENT

Let k be an algebraically closed field. We work in the category of algebraic varieties over k . $H^*(X)$ will denote étale cohomology with \mathbb{Q}_ℓ -coefficients for some ℓ prime to the characteristic of k . If $k = \mathbb{C}$, we take Betti cohomology with \mathbb{Q} -coefficients.

Let C be a smooth, complete curve over k , and let $\delta \subset C$ be a nonempty finite set of reduced points. Let $J(C)$ be the Jacobian of C , and let $J(C, \delta)$ be the semiabelian variety which represents the functor

$$(2.1) \quad X \mapsto \{(\mathcal{L}, \phi) \mid \mathcal{L} \text{ line bundle on } C \times X, \deg \mathcal{L}|_{C \times k(X)} = 0 \\ \phi : \mathcal{L}|_{\delta \times X} \cong \mathcal{O}_{\delta \times X}\} / \cong .$$

There is an exact sequence

$$(2.2) \quad 0 \rightarrow T \rightarrow J(C, \delta) \rightarrow J(C) \rightarrow 0$$

where T is the torus $\Gamma(\delta, \mathcal{O}^\times) / \Gamma(C, \mathcal{O}^\times)$. By abuse of notation we shall write $J(C, \delta)$ rather than $J(C, \delta)(k)$. We can identify the character group $\text{Hom}(T, \mathbb{G}_m)$ with $\text{Div}_\delta^0(C)$, the group of 0-cycles of degree 0 supported on δ . A split subgroup $\Delta \subset \text{Div}_\delta^0(C)$ corresponds to a quotient $T \twoheadrightarrow T_\Delta = T / \ker \Delta$, where $\ker \Delta \subset T$ is the subtorus killed by all characters in Δ . We may push out (2.2) and define $J(C, \Delta) := J(C, \delta) / \ker \Delta$:

$$(2.3) \quad 0 \rightarrow T_\Delta \rightarrow J(C, \Delta) \rightarrow J(C) \rightarrow 0.$$

The functor represented by $J(C, \Delta)$ is the following quotient of (2.1)

$$(2.4) \quad X \mapsto \{(\mathcal{L}, \phi) \mid \mathcal{L} \text{ line bundle on } X \times C, \deg \mathcal{L}|_{k(X) \times C} = 0 \\ \phi : \otimes_i \mathcal{L}^{\otimes n_i}|_{X \times \{c_i\}} \cong \mathcal{O}_X \text{ for all } \sum n_i c_i \in \Delta\}.$$

These trivializations should be compatible in an evident way with the group law on Δ .

Lemma 2.1. *We write $H^1(C, \delta) = H_c^1(C \setminus \delta)$. Define*

$$H^1(C, \Delta) := (H^1(C, \delta)/\Delta^\perp) \otimes \mathbb{Q}_\ell,$$

where

$$\Delta^\perp \otimes \mathbb{Q}_\ell \subset \mathbb{Q}_\ell[\delta]/\mathbb{Q}_\ell \subset H^1(C, \delta; \mathbb{Q}_\ell)$$

is perpendicular to $\Delta \subset \text{Div}_\delta^0(C)$ under the evident coordinatewise duality. The first Chern class $c_1(\mathcal{L}_\Delta)$ of the Poincaré bundle \mathcal{L}_Δ on $C \times J(C, \Delta)$ lies in $H^1(C, \Delta) \otimes H^1(J(C, \Delta))(1)$.

Proof. Let $I_\delta \subset \mathcal{O}_{J(C, \Delta) \times C}$ be the ideal of $J(C, \Delta) \times \delta$. Let $\pi : C \rightarrow C'$ be the singular curve obtained from C by gluing all the points of δ to a single point $\delta' \in C'$. Define $M_\Delta \subset (1 \times \pi)_*(\mathcal{O}_{J(C, \Delta) \times C}^\times)/k^\times$ to be the pullback as indicated:

$$(2.5) \quad \begin{array}{ccccccc} 0 & \rightarrow & (1 \times \pi)_*(1 + I_\delta) & \rightarrow & M_\Delta & \rightarrow & (\ker \Delta)_{J(C, \Delta) \times \delta'} \rightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \rightarrow & (1 \times \pi)_*(1 + I_\delta) & \rightarrow & \frac{(1 \times \pi)_* \mathcal{O}_{J(C, \Delta) \times C}^\times}{k^\times} & \rightarrow & \frac{(1 \times \pi)_* \mathcal{O}_{J(C, \Delta) \times \delta}^\times}{k^\times} \rightarrow 0. \end{array}$$

One gets a diagram of Kummer sequences of sheaves on $J(C, \Delta) \times C'$ (Here $j : C \setminus \delta \hookrightarrow C$)

$$(2.6) \quad \begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & (1 \times \pi)_*(1 \times j)_* \mu_{\ell^n} & \rightarrow & (1 \times \pi)_*(1 + I_\delta) & \xrightarrow{\ell^n} & (1 \times \pi)_*(1 + I_\delta) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & M_{\Delta, \ell^n} & \rightarrow & M_\Delta & \xrightarrow{\ell^n} & M_\Delta \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & ((\ker \Delta)_{J(C, \Delta) \times \delta'})_{\ell^n} & \rightarrow & (\ker \Delta)_{J(C, \Delta) \times \delta'} & \xrightarrow{\ell^n} & (\ker \Delta)_{J(C, \Delta) \times \delta'} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

We have $[\mathcal{L}_\Delta] \in H^1(J(C, \Delta) \times C', M_\Delta)$ and so by the Kummer coboundary, $c_1(\mathcal{L}_\Delta) \in \varprojlim_n H^2(J(C, \Delta) \times C', M_{\Delta, \ell^n})$. But $M_{\Delta, \ell^n} \cong \mathbb{Z}/\ell^n_{J(C, \Delta)} \boxtimes \psi_{\ell^n}$, where ψ_{ℓ^n} fits into an exact sequence of sheaves on C'

$$(2.7) \quad 0 \rightarrow \pi_* j_! \mu_{\ell^n} \rightarrow \psi_{\ell^n} \rightarrow (\ker \Delta)_{\delta', \ell^n} \rightarrow 0.$$

We can identify $\Delta^\perp \otimes \mu_{\ell^n}$ with $(\ker \Delta)_{\ell^n}$. the exact cohomology sequence from (2.7) yields

$$(2.8) \quad (\ker \Delta)_{\mu_{\ell^n}} \rightarrow H^1(C, \delta; \mu_{\ell^n}) \rightarrow H^1(C', \psi_{\ell^n}) \rightarrow 0.$$

Passing to the limit over n , it now follows that we may define $c_1(\mathcal{L}_\Delta) \in H^1(J(C, \Delta)) \otimes H^1(C, \Delta)(1)$ as in the statement of the lemma. \square

Lemma 2.2. *Suppose given a morphism $\rho : X \rightarrow J(C)$. Let Ξ be a Cartier divisor on $C \times X$ representing ρ . We assume Ξ is flat over C so we may define a correspondence $\Xi_* : \text{Div}(C) \rightarrow \text{Div}(X)$. Let $U \subset X$ be nonempty open in X . Then there exists a lifting $\rho_{U, \Delta} : U \rightarrow J(C, \Delta)$ of ρ if and only if $(\Xi|_{C \times U})_*(\Delta) \subset \text{Div}(U)$ consists of principal divisors. The set of such liftings is a torsor under $\text{Hom}(\Delta, \Gamma(U, \mathcal{O}_U^\times))$.*

Proof. Choose a basis $z_i = \sum_j n_{ij} c_j$ for the free abelian group Δ . Write $\mathcal{O}_{C \times X}(\Xi)_{z_i \times X} := \otimes_j \mathcal{O}_{C \times X}(\Xi)^{\otimes n_{ij}}|_{\{c_j\} \times X}$. The assumption that $\Xi_*(\Delta)$ consists of principal divisors is precisely the assumption that all the line bundles $\mathcal{O}_{C \times X}(\Xi)_{z_i \times X}|_U$ are trivial. The choice of the trivializations for a basis of Δ yields the choice of the desired lifting $\rho_{U, \Delta}$. \square

Lemma 2.3. *Assume X is a smooth variety, and let $\rho : X \rightarrow J(C)$ be as above. Suppose $U \subset X$ is a dense open set such that $\rho|_U$ admits a lifting $\rho_{U, \Delta} : U \rightarrow J(C, \Delta)$. Let $\text{Div}_{X \setminus U}^0(X)$ be the free abelian group on Cartier divisors supported on $X \setminus U$ which are homologous to 0 on X . Then we get a commutative diagram on cohomology*

$$(2.9) \quad \begin{array}{ccccccc} 0 & \rightarrow & H^1(J(C)) & \rightarrow & H^1(J(C, \Delta)) & \rightarrow & \Delta \otimes \mathbb{Q}_\ell(-1) \rightarrow 0 \\ & & \downarrow \rho^* & & \downarrow \rho_{U, \Delta}^* & & \downarrow a \\ 0 & \rightarrow & H^1(X) & \rightarrow & H^1(U) & \rightarrow & \text{Div}_{X \setminus U}^0(X) \otimes \mathbb{Q}_\ell(-1) \rightarrow 0. \end{array}$$

Proof. The left hand square is commutative by functoriality. That the cokernels on the top and bottom row are as indicated follows on the top row from the Leray spectral sequence for the projection $\pi : J(C, \Delta) \rightarrow J(C)$ and on the bottom from the localization sequence which may be written

$$(2.10) \quad 0 \rightarrow H^1(X) \rightarrow H^1(U) \rightarrow H_{X \setminus U}^2(X) \rightarrow H^2(X).$$

The identification $H_{X \setminus U}^2(X) \cong \text{Div}_{X \setminus U}(X) \otimes \mathbb{Q}_\ell(-1)$ is saying that by purity, the Gysin homomorphism is an isomorphism. \square

Remark 2.4. (i) Fixing $\rho_{U,\Delta}$ amounts to fixing trivializations of the restriction $\mathcal{O}_{C \times X}(\Xi)|_{z_i \times X|_U}$ as above. Such trivializations exhibit

$$\mathcal{O}_{C \times X}(\Xi)|_{z_i \times X} \cong \mathcal{O}_X(D_i)$$

for some divisor D_i with support on $X \setminus U$. The map labeled a in (2.9) sends $z_i \mapsto D_i$.

(ii) The diagram

$$(2.11) \quad \begin{array}{ccc} \Delta & \longrightarrow & J(C) \\ \downarrow a & & \downarrow \rho^* \\ \mathrm{Div}_{X \setminus U}^0(X) & \longrightarrow & \mathrm{Pic}^0(X) \end{array}$$

is commutative, where the horizontal arrows are cycle classes. Indeed, both a and ρ^* are defined by the divisor on $C \times X$. Note that a depends on the choice of $\rho_{U,\Delta}$ but only up to rational equivalence.

Now suppose X is smooth, projective, of dimension d . Let $U \subset X$ be a dense open subset. Write $X \setminus U = D \cup Z$ where $D \subset X$ is a divisor and $\mathrm{codim}(Z \subset X) \geq 2$. We have $H^1(X \setminus D) \cong H^1(U)$. Since we are interested in $H^1(U)$, we may assume $U = X \setminus D$ is the complement of a divisor.

Let $i : C \hookrightarrow X$ be a general linear space section of dimension 1, and let $\delta = C \cap D$. We may choose $\rho : X \rightarrow J(C)$ such that the composition

$$(2.12) \quad \mathrm{Pic}^0(X) \xrightarrow{i^*} J(C) \xrightarrow{\rho^*} \mathrm{Pic}^0(X)$$

is multiplication by an integer $N \neq 0$. Indeed, let H be a very ample line bundle so that C is the $(d-1)$ -fold product of general sections of H . Intersection with H yields an isogeny $\mathrm{Pic}^0(X) \rightarrow \mathrm{Alb}(X)$, which defines an inverse isogeny $\mathrm{Alb}(X) \rightarrow \mathrm{Pic}^0(X)$ of degree N . We pull back the Poincaré bundle from $J(C) \times J(C)$ to $C \times X$ via the composite map $C \times X \rightarrow J(C) \times \mathrm{Alb}(X) \rightarrow J(C) \times \mathrm{Pic}^0(X) \rightarrow J(C) \times J(C)$, where the first map is the cycle map, the second one is $1 \times$ isogeny, the third one is $1 \times$ restriction. We define $\mathcal{O}_{C \times X}(\Xi)$ to be the inverse image of the Poincaré bundle. The morphism $\rho : X \rightarrow J(C)$ is the correspondence $x \mapsto \mathcal{O}_{C \times X}(\Xi)|_{C \times \{x\}}$ and does not depend on the choice of the section Ξ .

Consider the diagram

$$(2.13) \quad \begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Q}_\ell[\delta]/\mathbb{Q}_\ell & \rightarrow & H_c^1(C \setminus \delta) & \rightarrow & H^1(C) & \rightarrow 0 \\ & & \downarrow b & & \downarrow i_* & & \downarrow i_* & \\ 0 & \rightarrow & \frac{H^{2d-2}(D)}{H^{2d-2}(X)}(d-1) & \rightarrow & H_c^{2d-1}(U)(d-1) & \rightarrow & H^{2d-1}(X)(d-1) & \end{array}$$

Here, the rows are long exact sequences associated to restriction to closed subsets, and the vertical arrows are Gysin maps. The map b can be described as follows. The \mathbb{Q}_ℓ -vector space $H^{2d-2}(D)(d-1)$ has basis the irreducible components of D , and $b(x)$ is the basis element $[D_x]$ associated to the unique component D_x of D containing x . We have dual exact sequences (defining $\text{Div}_D^0(X)$)

$$(2.14) \quad 0 \rightarrow \text{Div}_D^0(X) \rightarrow H_{2d-2}(D)(1-d) \rightarrow H^2(X)(1) \\ H^{2d-2}(X)(d-1) \rightarrow H^{2d-2}(D)(d-1) \rightarrow \frac{H^{2d-2}(D)}{H^{2d-2}(X)}(d-1) \rightarrow 0.$$

If we view $\mathbb{Q}_\ell[\delta]$ and $H^{2d-2}(D)(d-1)$ as endowed with symmetric pairings with orthonormal bases the points $x \in \delta$ and the cohomology classes of irreducible components $D_i \subset D$, then b is adjoint to the map $D_i \mapsto D_i \cdot \delta$. We conclude

Lemma 2.5. *Define $\text{Div}_D^0(X)$ to be the \mathbb{Q}_ℓ -vector space spanned by divisors on X supported on D and homologous to 0 on X . Define $\Delta \subset \text{Div}_\delta^0(C)$ to be the image of $\text{Div}_D^0(X)$ under pullback i^* . Then there is a commutative diagram*

$$(2.15) \quad \begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Q}_\ell[\delta]/\Delta^\perp & \rightarrow & H_c^1(C, \Delta) & \rightarrow & H^1(C) & \rightarrow 0 \\ & & \downarrow b & & \downarrow i_* & & \downarrow i_* & \\ 0 & \rightarrow & \frac{H^{2d-2}(D)}{H^{2d-2}(X)}(d-1) & \rightarrow & H_c^{2d-1}(U)(d-1) & \rightarrow & H^{2d-1}(X)(d-1). \end{array}$$

Proof. The map b is dual to the restriction map $\text{Div}_D^0(X) \xrightarrow{i^*} \text{Div}_\delta^0(C)$. By definition Δ^\perp is orthogonal to the image of i^* , i.e. $\Delta^\perp = \ker b$. \square

Lemma 2.6. *Let $\Delta = i^*(\text{Div}_D^0(X)) \subset \text{Div}_\delta^0(C)$ be as in Lemma 2.5. Then ρ defined in (2.12) lifts to some $\rho_{U, \Delta} : U \rightarrow J(C, \Delta)$.*

Proof. The correspondence defined by $\mathcal{O}_{C \times X}(\Xi)$ in (2.12) carries $\mathcal{O}_C(z)$ for $z \in \Delta = i^*(\text{Div}_D^0(X))$ to line bundles in $\text{Pic}^0(X)$, the classes of which fall in the image of $\rho^* i^*(\text{Div}_D^0(X)) \cong N \cdot \text{Div}_D^0(X)$ in $\text{Pic}^0(X)$. To be more precise, let D_p be a basis for $\text{Div}_D^0(X)$, and set $z_p = i^* D_p$. This is a basis of Δ . Then $\mathcal{O}_{C \times X}(\Xi)|_{z_p \times X} = \mathcal{O}_X(D_p)$. Thus choose $\rho_{U, \Delta}$ in Lemma 2.2 using this trivialization on U . \square

Using the Lemmas 2.1, 2.5, 2.6 together with (2.9), we pull back

$$(2.16) \quad c_1(\mathcal{L}_\Delta) \in H^1(C, \Delta) \otimes H^1(J(C, \Delta))(1) \xrightarrow{i_* \otimes \rho_{U, \Delta}} \\ H_c^{2d-1}(U)(d) \otimes H^1(U) \cong H^1(U)^\vee \otimes H^1(U)$$

and define a correspondence $\Phi : H^1(U) \rightarrow H^1(U)$.

Lemma 2.7. *The map Φ is the multiplication by N .*

Proof. We consider Φ . It acts on $H^1(U)$, compatibly with the exact sequence

$$(2.17) \quad 0 \rightarrow H^1(X) \rightarrow H^1(U) \rightarrow \mathrm{Div}_D^0(X)(-1) \rightarrow 0$$

By definition of $\rho_{U,\Delta}$, it is equal to $N \cdot \mathrm{Id}$ on $H^1(X)$ and on $\mathrm{Div}_D^0(X)(-1)$. Thus $\Phi - N \cdot \mathrm{Id}$ is a correspondence from $\mathrm{Div}_D^0(X)(-1)$ to $H^1(X)$. We use purity in the sense of Deligne. There is no nontrivial correspondence $\mathrm{Div}_D^0(X)(-1) \rightarrow H^1(X)$. If $k = \mathbb{C}$ and we consider Betti cohomology, $\mathrm{Div}_D^0(X)(-1)$ is pure of weight 2 while $H^1(X)$ is pure of weight 1. If k is the algebraic closure of a finite field, we have the same conclusion. Otherwise, all the objects used are defined over a finitely generated field k over a finite field k_0 . By Chebotarev theorem, the Galois group of k/k_0 is generated by Frobenii, so we may make sense of the notion of weight for $H^1(U)$. We conclude as in the complex case. \square

We now express in terms of cycles the trivialization of $\mathcal{O}_{C \times X}(\Xi)_{p \times X} = \mathcal{O}_X(D_p)$ used in the proof of Lemma 2.6.

Theorem 2.8. *With notation as above, there exists a cycle Γ on $X \times U$ of dimension $d = \dim X$ together with rational functions f_μ on X for each divisor μ homologically equivalent to 0 on X and supported on $D = X_U$ such that $pr_{2*}(\Gamma \cdot (\mu \times U)) = (f_\mu)$. The data $(\Gamma, \{f_\mu\})$ define a class in $H_c^{2d-1}(U) \otimes H^1(U)$ which gives the identity map on $H^1(U)$.*

We close this section with a comment about Künneth projectors $\pi^i : H^*(U) \rightarrow H^i(U)$ for $i > 1$. We consider the somewhat weaker question of the existence of an algebraic projector when we localize at the generic point of the target, i.e. we consider $H^*(U) \rightarrow H^i(U) \rightarrow \varprojlim_{V \subset U} H^i(V)$. We assume $U = X \setminus D$ with X smooth, projective, and D a Cartier divisor.

Proposition 2.9. *Let $n < \dim X$ be an integer. Let $Y \subset X$ be a plane section of dimension n which is general with respect to D . Write $\delta = Y \cap D$. Then the restriction map*

$$(2.18) \quad H_D^{n+1}(X) \rightarrow H_\delta^{n+1}(Y)$$

is injective.

Proof. Let $d = \dim(X)$. By duality, we have to show surjectivity of the Gysin map $H^{n-1}(\delta) \rightarrow H^{2d-(n+1)}(D)(d-n)$. More generally, one has

Theorem 2.10 (P. Deligne). *Let \mathcal{F} be a ℓ -adic sheaf on \mathbb{P}^n , then for A in a non-trivial open subset of the dual projective space $(\mathbb{P}^n)^\vee$, the Gysin homomorphism*

$$H^{i-2}(A, \iota^* \mathcal{F})(-1) \rightarrow H_A^i(\mathbb{P}^n, \mathcal{F}),$$

where $\iota : A \rightarrow \mathbb{P}^n$ is the closed embedding, is an isomorphism for all i . In particular, if $V \subset \mathbb{P}^n$ is a projective variety, then the Gysin homomorphism $H^i(A \cap V) \rightarrow H^{i+2}(V)(1)$ is an isomorphism for $i > \dim(A \cap V)$ and surjective for $i = \dim(A \cap V)$ for A in a nonempty open subset of $(\mathbb{P}^n)^\vee$.

The proof of the general theorem is written in [5], Theorem 2.1. Applied to $\mathcal{F} = a_* \mathbb{Q}_\ell$, where $a : V \rightarrow \mathbb{P}^n$ is the projective embedding, it shows that the Gysin isomorphism $H^i(A \cap V) \rightarrow H_{A \cap V}^{i+2}(V)(1)$ is an isomorphism. Then the application follows from Artin's vanishing theorem $H^i(V \setminus (A \cap V)) = 0$ for $i > \dim(V)$. \square

Let L be the Lefschetz operator on $H^*(X)$. One of the standard conjectures ($B(X, L)$ in [10]) is the existence of an algebraic correspondence Λ which is a "weak inverse" to L . Assume now that this standard conjecture B is true for X and for all smooth linear space sections $Y \subset X$. The strong Lefschetz theorem implies that $L^{d-n} : H^n(X) \xrightarrow{\cong} H^{2d-n}(X)(d-n)$. Assuming $B(X, L)$, $\Lambda^{d-n} = (L^{d-n})^{-1} : H^{2d-n}(X)(d-n) \cong H^n(X)$. Write $P = \Lambda^{d-n}|_{Y \times X}$. It is easy to check that the composition

$$(2.19) \quad H^n(X) \xrightarrow{i^*} H^n(Y) \xrightarrow{P} H^n(X)$$

is the identity, so $H^n(Y) = \text{Image}(i^*) \oplus \ker(P)$. Consider the diagram

$$(2.20) \quad \begin{array}{ccccc} H^n(X) & \rightarrow & H^n(U) & \xrightarrow{a} & H_D^{n+1}(X) \\ \downarrow i^* & & \downarrow & & \downarrow b \\ H^n(Y) & \xrightarrow{d} & H^n(Y \setminus \delta) & \xrightarrow{c} & H_\delta^{n+1}(Y) \end{array}$$

Define

$$(2.21) \quad H^n(Y \setminus \delta)^0 = \{x \in H^n(Y \setminus \delta) \mid c(x) \in \text{Im}(b \circ a)\}.$$

As a consequence of proposition 2.9 and (2.20) we see that $H^n(U) \rightarrow H^n(Y \setminus \delta)^0/d(\ker(P))$, and the kernel of this map is the image in $H^n(U)$ of elements $x \in H^n(X)$ such that $i^*x \in \ker(P) \oplus \text{Image}(H_\delta^n(Y) \rightarrow H^n(Y))$. For such an x , it will necessarily be the case that $x = P(i^*x)$

is supported on a proper closed subset of X . In particular, for some $V \subset U$ open dense, P will induce a map

$$(2.22) \quad P_U : H^n(Y \setminus \delta)^0 \rightarrow H^n(V).$$

The map $i^* : H^n(U) \rightarrow H^n(Y \setminus \delta)^0$ dualizes to $i_* : (H^n(Y \setminus \delta)^0)^\vee \rightarrow H_c^{2d-n}(U)$, so we may define

$$(2.23) \quad (i_* \otimes P_U) : H^n(Y \setminus \delta)^0{}^\vee \otimes H^n(Y \setminus \delta)^0 \rightarrow H_c^{2d-n}(U) \otimes H^n(V).$$

Let $T \subset H^n(Y \setminus \delta)^0$ be the subgroup of cohomology classes supported in codimension 1. Assuming inductively that we are able to define an algebraic correspondence on Y which carries a class

$$(2.24) \quad \gamma \in (H^n(Y \setminus \delta)^0)^\vee \otimes (H^n(Y \setminus \delta)^0/T)$$

corresponding to the evident map $H^n(Y \setminus \delta)^0 \twoheadrightarrow H^n(Y \setminus \delta)^0/T$, it would follow since $P_U(T) \subset \ker(H^n(V) \rightarrow \varprojlim_{V \subset U} H^n(V))$ that we could view

$$(2.25) \quad (i_* \otimes P_U)(\gamma) \in H_c^{2d-n}(U) \otimes \varinjlim_{V \subset U} H^n(V).$$

This correspondence would have the desired properties.

One interest in pursuing this line of investigation concerns the Milnor conjecture that the Galois cohomology with $\mathbb{Z}/n\mathbb{Z}$ -coefficients prime to the residue characteristic is generated as an algebra by H^1 . There is a geometric proof of this result in top degree [2], so, for example, elements in $H^n(Y \setminus \delta)$ lie in the subalgebra generated by H^1 after localization. If P_U exists as an algebraic correspondence, then using the existence of a norm in Milnor K -theory, one could show that the Milnor conjecture was true for $H^*(k(X), \mathbb{Z}/\ell\mathbb{Z})$ for almost all ℓ . (The condition on ℓ arises because the standard conjectures only make sense after tensoring with \mathbb{Q} .) Here, the idea that cohomology classes in degree n might come by correspondence from an algebraic variety of dimension $\leq n$ was suggested to us by A. Beilinson.

3. OPEN CORRESPONDENCES

The aim of this section is to give a simple motivic proof of the independence of ℓ or of a complex embedding of a ground field k of the trace for open correspondences. If we assume that k is finite, then, as conjectured by Deligne, high Frobenius power twists move the correspondence to a general position correspondence and the local factors have been computed in [3], [13], [14], [6]. Surely in this case the simple observations which follow are weaker.

We consider open correspondences. This means the following. Let X be a smooth projective variety of dim d over an algebraically closed field k , and let $U \subset X$ be a nontrivial open subvariety, with complement

$D = X \setminus U$. One considers codim d cycles $\Gamma \subset U \times U$ which have the property that they induce a correspondence $\Gamma_* : H^i(U) \rightarrow H^i(U)$ or equivalently $H_c^i(U) \rightarrow H_c^i(U)$ for all i . Here cohomology is étale \mathbb{Q}_ℓ cohomology or Betti cohomology if $k = \mathbb{C}$ and we denote by $p_i : X \times X \rightarrow X$ the two projections. We write

$$(3.1) \quad \Gamma = \sum n_j \Gamma_j$$

where Γ_j is irreducible, $n_j \in \mathbb{Z}$ and define

$$(3.2) \quad \bar{\Gamma} := \sum n_j \bar{\Gamma}_j, \quad \Gamma_j \subset U \times U$$

where $\bar{}$ is the Zariski closure in $X \times X$.

Definition 3.1. If $p_{2|(\Gamma)_j} : \Gamma_j \rightarrow U$ is proper for all j , or equivalently if

$$(3.3) \quad \bar{\Gamma}_j \cap (D \times X) \subset X \times D \quad \forall j,$$

then one defines

$$(3.4) \quad (\Gamma_j)_* : H_c^i(U) \xrightarrow{p_{2*}} H_c^i(\Gamma_j) \xrightarrow{(p_1)_*} H_c^i(U),$$

and call it the open correspondence defined by Γ_j . The correspondence defined by Γ is then by definition $\Gamma_* = \sum n_j (\Gamma_j)_*$.

Remark 3.2. If we compare this condition to the one yielding to Künneth correspondences in sections 2 and 3, then it is much stronger. Indeed, we had $P \subset W \times X$, thus $\bar{\Gamma} = P$ viewed as a cycle in $X \times X$ and $P \cap (D \times X) = P \cap (\delta \times X)$ was not empty.

We assume now that we have the assumption as in Definition 3.1 and we wish to give conditions under which one can compute the trace of Γ_* which is defined by

$$(3.5) \quad \text{Tr}(\Gamma_*) := \sum_{i=0}^{2d} (-1)^i \text{Tr}(\Gamma_* |_{H_c^i(U)}).$$

As it stands, the trace of Γ_* depends a priori on ℓ , or, for varieties defined over a field k of characteristic 0, and Betti cohomology taken with respect to a complex embedding $\iota : k \rightarrow \mathbb{C}$, it depends on ι . One has

Theorem 3.3. *Let X be a smooth projective variety of dimension d defined over a field k , together with a strict normal crossings divisor $D \subset X$ of open complement $U = X \setminus D$. Let $\Gamma \subset U \times U$ be a dimension d cycle defining an open correspondence Γ_* on ℓ -adic cohomology or Betti cohomology as in Definition 3.1. Then $\text{Tr}(\Gamma_*)$ does not depend*

on ℓ in ℓ -adic cohomology or on the complex embedding of k in Betti cohomology.

Proof. We use the relative motivic cohomology $H_M^{2d}(X \times U, D \times U, \mathbb{Z}(d))$, as defined in [11], chapter 4, 2.2 and p. 209. The group $H_M^m(X \times U, D \times U, \mathbb{Z}(n))$ is the homology $H_{2n-m}(\mathcal{Z}^n(X \times U, D \times U, *))$, where $\mathcal{Z}^n(X \times U, D \times U, *)$ is the single complex associated to the double higher Chow cycle complex

$$(3.6) \quad \begin{array}{ccccc} \dots & & \dots & & \dots \\ \partial \downarrow & & \partial \downarrow & & \downarrow \\ \mathcal{Z}^n(X \times U, 1) & \xrightarrow{\text{rest}} & \mathcal{Z}^n(D^{(1)} \times U, 1) & \xrightarrow{\text{rest}} & \mathcal{Z}^n(D^{(2)} \times U, 1) \\ \partial \downarrow & & \partial \downarrow & & \downarrow \\ \mathcal{Z}^n(X \times U, 0) & \xrightarrow{\text{rest}} & \mathcal{Z}^n(D^{(1)} \times U, 0) & \xrightarrow{\text{rest}} & \mathcal{Z}^n(D^{(2)} \times U, 0). \end{array}$$

Here $D^{(a)}$ is the normalization of all the strata of codimension a , $\mathcal{Z}^n(D^{(a)} \times U, b)$ is a group of cycles on $D^{(a)} \times U \times S^b$ where S^\bullet is the cosimplicial scheme $S^n = \text{Spec}(k[t_0, \dots, t_n]/(\sum_{i=0}^n t_i - 1))$ with face maps $S^n \hookrightarrow S^{n+1}$ defined by $t_i = 0$. More precisely, $\mathcal{Z}^n(D^{(a)} \times U, b)$ is generated by the codimension n subvarieties $Z \subset D^{(a)} \times U \times S^b$ such that, for each face F of S^b , and each irreducible component $F' \subset D^{(a)}$ of the strata of D we have $\text{codim}_{F' \times U \times F}(Z \cap (F' \times U \times F)) \geq n$. The horizontal restriction maps are the intersection with the smaller strata, the vertical ∂ 's are the boundary maps.

This relative motivic cohomology acts as correspondences on $H_c^*(U)$, where $H_c^*(U)$ is ℓ -adic or Betti cohomology.

Let us write $\Gamma = \sum n_j \Gamma_j$. By (3.3), one has $\Gamma_j \subset X \times U$ closed with $\Gamma_j \cap (D \times U) = \emptyset$, thus in particular, $\Gamma \in \mathcal{Z}^d(X \times U, 0)$ with $\text{rest}(\Gamma) = 0$ in $\mathcal{Z}^d(D^{(1)} \times U, 0)$, thus it defines a class

$$(3.7) \quad [\Gamma] \in H^{2d}(X \times U, D \times U, d).$$

Similarly, we consider the restriction $\Delta_U \subset U \times X$ of the diagonal $\Delta \subset X \times X$. This defines a class in $\mathcal{Z}^d(U \times X, 0)$. As $\text{rest}(\Gamma_U) = 0$ in $\mathcal{Z}^d(U \times D^{(1)}, 0)$, it defines a class

$$(3.8) \quad [\Delta_U] \in H^{2d}(U \times X, U \times D, d).$$

We want to pair $[\Gamma]$ with $[\Delta_U]$. We argue using M. Levine's work. Let Y be a N -dimensional smooth projective defined over k , with two strict normal crossings divisors A, B so that $A+B$ is a strict normal crossings divisor. By [11], Chapter IV, lemma 2.3.5 and lemma 2.3.6, the motive

$M(Y \setminus A, B \setminus B \cap A)$ is dual to the motive $M(Y \setminus B, A \setminus B \cap A)$. It yields a cup product

$$(3.9) \quad H_M^a(Y \setminus A, B \setminus B \cap A, b) \times H_M^{2N-a}(Y \setminus B, A \setminus A \cap B, N-b) \rightarrow \mathbb{Z}.$$

This cup product is compatible with the cup product in ℓ -adic or Betti cohomology. We apply this to

$$(3.10) \quad Y = X \times X, \quad A = D \times X, \quad B = X \times D$$

so we can cup $[\Delta_U] \in H_M^{2d}(Y \setminus A, B \setminus B \cap A, d)$ and $[\Gamma] \in H_M^{2d}(Y \setminus B, A \setminus A \cap B, d)$

$$(3.11) \quad [\Delta_U] \cup [\Gamma] \in \mathbb{Z}.$$

The theorem is then the consequence of the following proposition.

Proposition 3.4.

$$\mathrm{Tr}(\Gamma_*) = [\Delta_U] \cup [\Gamma] \in \mathbb{Z}.$$

Proof. By the compatibility of the cup product (3.9) with cohomology, we just have to prove the proposition with $[\Delta_U]$ and $[\Gamma]$ replaced by their classes $\mathrm{cl}(\Delta_U)$ and $\mathrm{cl}(\Gamma)$ in cohomology. We may assume that $\Gamma \subset X \times U$ is irreducible. On the other hand, the map

$$(3.12) \quad H_c^i(\Gamma) \xrightarrow{p_{1*}} H_c^i(U)$$

factors through

$$(3.13) \quad \begin{aligned} H_c^i(\Gamma) &\xrightarrow{\mathrm{Gysin}} H_{\Gamma, c}^{i+2d}(X \times U, d) = H_{\Gamma, c}^{i+2d}(U \times U, d) \\ &\rightarrow H_c^{i+2d}(U \times U, d) \rightarrow H_c^{i+2d}(U \times X, d) \xrightarrow{p_{1*}} H_c^i(U) \end{aligned}$$

so the correspondence

$$(3.14) \quad \Gamma_* : H_c^i(U) \xrightarrow{p_2^*} H_c^i(\Gamma) \xrightarrow{p_{1*}} H_c^i(U)$$

is just

$$(3.15) \quad H_c^i(U) \cup (\mathrm{cl}(\Gamma) \in H^{2d}(X \times U, D \times U, d)) \subset H_c^{i+2d}(U \times X, d)$$

followed by p_{1*} . Now we argue as in the classical case. Let e_a^i be a basis of $H_c^i(U)$, and $(e_a^i)^\vee$ be its dual basis in $H^{2d-i}(U)(d)$. Write $\mathrm{cl}(\Gamma) = \sum_i \sum_a f_a^i \otimes (e_a^i)^\vee$, $f_a^i = \sum f_{ab}^i e_b^i \in H_c^i(U)$. So $\Gamma_*(e_a^i) = \sum_b f_{ab}^i$, and $\mathrm{Tr}(\Gamma_*) = \sum (-1)^i \sum_a \sum_b f_{ab}^i$. On the other hand, one has $\mathrm{cl}(\Delta_U) = \sum_i \sum_a (e_a^i)^\vee \otimes (e_a^i)$. Thus $\mathrm{cl}(\Delta_U \cup \Gamma) = \sum (-1)^i \sum_a \sum_b f_{ab}^i$. □

The proposition finishes the proof of the theorem. □

The rest of the section is devoted to giving a concrete expression for (3.5) under stronger geometric assumptions on Γ .

The condition (3.3) allows to define the embedding

$$(3.16) \quad \begin{aligned} \Gamma_j &= (p_2|_{\Gamma_j})^{-1}(U) = \bar{\Gamma}_j \setminus \bar{\Gamma}_j \cdot (X \times D) \\ &\xrightarrow{\iota \subset} (\Gamma_j)' := (p_1|_{\Gamma_j})^{-1}(U) = \bar{\Gamma}_j \setminus \bar{\Gamma}_j \cdot (D \times X). \end{aligned}$$

By definition

$$(3.17) \quad p_1|_{(\Gamma_j)'} \text{ is proper.}$$

Lemma 3.5. *For all j , one has a factorization*

$$(\Gamma_j)_* : H_c^i(U) \xrightarrow{p_2^*} H_c^i(\Gamma_j) \xrightarrow{\iota_*} H_c^i((\Gamma_j)') \xrightarrow{(p_1)_*} H_c^i(U).$$

Proof. By definition one has a factorization

$$(3.18) \quad \begin{array}{ccccc} \Gamma_j & \xrightarrow{\iota} & (\Gamma_j)' & \xrightarrow{\mu} & \bar{\Gamma}_j \\ p_1|_{\Gamma_j} \downarrow & & \downarrow p_1|_{(\Gamma_j)'} & & \downarrow p_1|_{\bar{\Gamma}_j} \\ U & \xrightarrow{=} & U & \xrightarrow{j} & X \end{array}$$

Setting $\lambda = \mu \circ \iota$, the lemma then just says that on p_1 -acyclic locally constant sheaves, one has a factorization $(p_1|_{\bar{\Gamma}_j})_! \lambda_! \rightarrow (p_1|_{\bar{\Gamma}_j})_! \mu_! \xrightarrow{\text{Tr}} j_!$. \square

Corollary 3.6. *One has a commutative diagram*

$$\begin{array}{ccc} H_c^i(U) & \rightarrow & H^i(X) \\ p_2^* \downarrow & & p_2^* \downarrow \\ H_c^i(\Gamma_j) & \rightarrow & H^i(\bar{\Gamma}_j) \\ \iota_* \downarrow & & = \downarrow \\ H_c^i((\Gamma_j)') & \rightarrow & H^i(\bar{\Gamma}_j) \\ p_1_* \downarrow & & p_1_* \downarrow \\ H_c^i(U) & \rightarrow & H^i(X) \end{array}$$

Proof. With the notations of (3.18), the commutativity of the two upper squares simply means on constant sheaves $p_2 j_! = \lambda_! p_2^* \rightarrow \mu_! p_2^* = \mu_!$. Thus the assertion is on the lower square. By (3.17), one has $R((p_1)_*)|_{\Gamma_j'} = R((p_1)_!)|_{\Gamma_j'}$ thus $R((p_1)_*)|_{\bar{\Gamma}_j} \mu_! = R((p_1)_!)|_{\bar{\Gamma}_j} \mu_! = j_! = R((p_1)_!)|_{\Gamma_j'} = j_! R((p_1)_*)|_{\Gamma_j'}$. This shows the commutativity. \square

We remark that in Corollary 3.6, the Grothendieck-Lefschetz trace formula allows to compute the trace of the correspondence $\bar{\Gamma}_*$ on X

$$(3.19) \quad \mathrm{Tr}(\bar{\Gamma}_*) = \deg(\bar{\Gamma} \cdot \Delta_X)$$

Thus, in the corollary, we would like to complete the commutative diagram in an exact sequence of commutative diagrams, so that we can apply the trace formula on all the terms but the one we seek. In the sequel, we give a strong geometric condition under which it is possible.

Definition 3.7. We assume that D is a strict normal crossings divisor. The dim d cycle $\Gamma \subset U \times U$ is said to be in good position with respect to $D \times X$ if the following two conditions are fulfilled.

- i) Each $\bar{\Gamma}_j$ cuts each stratum $D_I \times X$ in $\mathrm{codim} \geq d$, where $D_I = D_{i_1} \cap \dots \cap D_{i_r}$ for $I = \{i_1, \dots, i_r\}$ with $|I| = r$.
- ii)

$$\bar{\Gamma}_j \cap (D_I \times X) \subset D_I \times D_I$$

set theoretically.

In this case, for all I we define the cycles

$$(3.20) \quad Z_{jI} = \bar{\Gamma}_j \cdot (D_I \times X) \subset D_I \times D_I$$

Let us be more precise. We denote motivic cohomology by $\mathcal{H}^a(b)$. We drop the subscript j , thus $\Gamma = \Gamma_j$. This defines

$$(3.21) \quad Z_I = \sum m_{I,a} Z_{I,a}$$

as in (3.10), where the $Z_{I,a}$ are the reduced irreducible components of Z_I . One has the Gysin isomorphisms

$$(3.22) \quad \bigoplus_a \mathbb{Q}_\ell[Z_{I,a}] \xrightarrow{\cong} H_{Z_I}^{2(d-r)}(D_I \times D_I, d-r) \xrightarrow{\cong} H_{Z_I}^{2d}(D_I \times X, d)$$

This yields the commutative diagram

$$(3.23) \quad \begin{array}{ccc} \bigoplus_a \mathbb{Q} \cdot [Z_{I,a}] & \xrightarrow{\otimes_{\mathbb{Q}} \mathbb{Q}_\ell} & \bigoplus_a \mathbb{Q}_\ell \cdot [Z_{I,a}] \\ \cong \downarrow & & \downarrow \cong \\ \mathcal{H}_{Z_I}^{2(d-r)}(D_I \times D_I, d-r) & \xrightarrow{\otimes_{\mathbb{Q}} \mathbb{Q}_\ell} & H_{Z_I}^{2(d-r)}(D_I \times D_I, d-r) \\ \cong \downarrow & & \downarrow \cong \\ \mathcal{H}_{Z_I}^{2d}(D_I \times X, d) & \xrightarrow{\otimes_{\mathbb{Q}} \mathbb{Q}_\ell} & H_{Z_I}^{2d}(D_I \times X, d) \end{array}$$

So we conclude that Z_I is a well defined cycle

$$(3.24) \quad Z_I \in \mathcal{H}_{Z_I}^{2(d-r)}(D_I \times D_I, d-r) \rightarrow \mathcal{H}^{2(d-r)}(D_I \times D_I, d-r),$$

which defines a correspondence

$$(3.25) \quad (Z_I)_* = \sum_a m_{I,a} (Z_{I,a})_* : H^i(D_I) \rightarrow H^i(D_I),$$

by the formula

$$(3.26) \quad (Z_{I,a})_* : H^i(D_I) \xrightarrow{p_2^*} H^i(Z_{I,a}) \xrightarrow{(p_1)_*} H^i(D_I).$$

It can be described as follows. Let $(Z_{I,a})_i \in H^{2d-2r-i}(D_I) \otimes H^i(D_I)(d-r)$ be the Künneth component of $Z_{I,a}$. Then one has

$$(3.27) \quad (Z_{I,a})_* : H^i(D_I) \xrightarrow{p_2^*} H^i(D_I) \otimes H^0(D_I) \xrightarrow{\cup(Z_{I,a})_i} \\ H^i(D_I) \cup H^{2d-2r-i}(D_I)(d-r) \otimes H^i(D_I) \xrightarrow{\text{Tr} \otimes 1} H^i(D_I).$$

We denote by $\Delta_I \subset D_I \times D_I$ the diagonal. By the Grothendieck-Lefschetz trace formula, one has

$$(3.28) \quad \text{Tr}((Z_I)_*) = \deg(Z_I \cdot \Delta_I).$$

Lemma 3.8. *Let $J \subset I$ with $|J| = s, |I| = r$. One has a commutative diagram*

$$\begin{array}{ccc} H^i(D_I) & \xrightarrow{\text{rest}} & H^i(D_J) \\ (Z_I)_* \downarrow & & \downarrow (Z_J)_* \\ H^i(D_J) & \xrightarrow{\text{rest}} & H^i(D_J) \end{array}$$

Proof. We have $Z_I = \bar{\Gamma} \cdot (D_I \times X), Z_J = \bar{\Gamma} \cdot (D_J \times X) = Z_I \cdot (D_J \times X)$. One has the diagram

$$(3.29) \quad \begin{array}{ccc} Z_J & \xrightarrow{\subset} & D_J \times D_J \\ & & \downarrow \subset \\ D_J \times D_I & \xrightarrow{\subset} & D_J \times D_I \\ & & \uparrow \subset \\ & & Z_I \end{array}$$

yielding the commutative diagram

$$(3.30) \quad \begin{array}{ccc} H^{2d-2r-i}(D_I) \otimes H^i(D_I)(d-r) & \xrightarrow{1 \otimes \text{rest}} & H^{2d-2r-i}(D_I) \otimes H^i(D_J)(d-r) \\ & & \uparrow \text{Gysin} \otimes 1 \\ & & H^{2d-2s-i}(D_J) \otimes H^i(D_J)(d-s) \end{array}$$

Via this diagram, one has

$$(3.31) \quad (1 \otimes \text{rest})((Z_I)_i) = (\text{Gysin} \otimes 1)((Z_J)_i)$$

Here the notation Z_I means the sum aover all components with multiplicities, and similarly for the Künneth components. Thus we apply the description (3.27) to conclude. \square

In order to have a unified notation, we set for $|I| = 0$

$$(3.32) \quad Z_I = \bar{\Gamma}, \quad \Delta_I = \Delta_X.$$

One has

Theorem 3.9. *Let X be a smooth projective variety over an algebraically closed field k , $D \subset X$ be a strict normal crossing divisor, with complement $U = X \setminus D$, and $\Gamma \subset U \times U$ be a dim d correspondence, with $p_2|_{\Gamma} : \Gamma \rightarrow U$ proper, and in good position with respect to $D \times X$ in the sense of Definition 3.7. Then one has*

$$\text{Tr}(\Gamma_*) = \sum_{r=0}^d (-1)^r \sum_{|I|=r} \text{deg}(Z_I \cdot \Delta_I).$$

Proof. One has the Mayer-Vietoris spectral sequence

$$(3.33) \quad E_1^{a,b} = \bigoplus_{|I|=a} H^b(D_I) \Rightarrow H_c^{a+b}(U)$$

On the other hand, the correspondence Γ_* on $H_c^\bullet(U)$ is compatible with the correspondence $\bar{\Gamma}_*$ on $H^\bullet(X)$ by Corollary 3.6, and with the correspondence $(Z_I)_*$ in $H^\bullet(D_I)$ by Lemma 3.8. Thus Γ_* acts on the whole spectral sequence via $\bar{\Gamma}_*$, $(Z_I)_*$. One obtains

$$(3.34) \quad \text{Tr}(\Gamma_*) = \sum_{r=0}^d (-1)^r \sum_{|I|=r} \text{Tr}((Z_I)_*).$$

We conclude with (3.19), (3.28). \square

We now give a scheme-theoretic condition under which the expression given in Theorem 3.9 depends only on local contributions in U . This condition is inspired by [8], Lemma 2.3.1.

Definition 3.10. We assume that D is a strict normal crossings divisor. The dim d cycle $\Gamma \subset U \times U$ is said to be in scheme theoretic good position with respect to $D \times X$ if it is in good position in the sense of Definition 3.7 and ii) is replaced by

ii')

$$\bar{\Gamma}_j \cap (D_I \times X) \subset D_I \times D_I$$

scheme theoretically.

Proposition 3.11. *Let X be a smooth projective variety over an algebraically closed field k , $D \subset X$ be a strict normal crossing divisor, with complement $U = X \setminus D$, and $\Gamma \subset U \times U$ be a $\dim d$ correspondence, with $p_2|_\Gamma : \Gamma \rightarrow U$ proper, and in scheme theoretic good position with respect to $D \times X$ in the sense of Definition 3.10. We assume moreover that $\bar{\Gamma}_j$ and $\Delta|_U$ cut transversally. Then $\text{Tr}(\Gamma_*) = \deg(\Delta_U \cdot \Gamma)$.*

Proof. Due to the good position assumption, all intersection multiplicities are 1 and the contributions lying on $(D \times X) \cup (X \times D)$ cancel in Theorem 3.9. □

Example 3.12. One case where the conditions of proposition 3.11 hold is in characteristic p when Γ is the graph of Frobenius. In this case, of course, the result is known by other methods.

Example 3.13. This example is inspired by [8], Remark 2.3.6. We take $X = \mathbb{P}^1$, $D = \{\infty\}$, $U = \mathbb{A}^1$, $\Gamma = \Gamma_{pq} = \{x^p - y^q = 0\} \subset \mathbb{A}^1 \times \mathbb{A}^1$. Then Γ defines an open correspondence with $\text{Tr}(\Gamma_*) = p$. On the other hand, one has

$$(3.35) \quad \deg(\Gamma_{pq} \cdot \Delta_U) = \dim k[t]/(t^p - t^q) = \begin{cases} \max(p, q) & \text{if } p \neq q \\ \infty & \text{if } p = q \end{cases}$$

and one has

$$(3.36) \quad \deg(\bar{\Gamma}_{pq} \cdot (\infty \times \mathbb{P}^1)) = q, \quad \deg(\bar{\Gamma}_{pq} \cdot (\mathbb{P}^1 \times \infty)) = p.$$

Thus Γ_{pq} is in scheme theoretic good position with respect to $\infty \times \mathbb{P}^1$ if and only if $p > q$, and is always in good position with respect to $\infty \times \mathbb{P}^1$. Since $\deg(\Delta_{\mathbb{P}^1} \cdot \bar{\Gamma}_{pq}) = \deg(\mathcal{O}(1, 1) \cdot \mathcal{O}(p, q)) = p + q$, we see exactly how the formula of Theorem 3.9 works both in Theorem 3.9 and in Proposition 3.11.

REFERENCES

- [1] Beilinson, A.: Higher regulators and values of L -functions, *J. Soviet Math.* **30** (1985), 2036-2070.
- [2] Bloch, S.: Lecture on Algebraic cycles, *Duke University Mathematics Series*, IV, (1980).
- [3] Deligne: Rapport sur la formule des traces, SGA 4,5, *Lecture Notes in Mathematics* **569**, Springer Verlag.
- [4] Deligne, P.: Théorèmes de finitude en cohomologie ℓ -adique, in SGA 4 $\frac{1}{2}$, *Lect. Notes Math.* vol. **569**, 233-251, Berlin Heidelberg New York Springer 1977.
- [5] Esnault, H.: Eigenvalues of Frobenius acting on the ℓ -adic cohomology of complete intersections of low degree, *C. R. Acad. Sci. Paris, Ser. I* **337** (2003), 317-320.
- [6] Fujiwara, K.: Rigid geometry, Lefschetz-Verdier trace formula and Deligne's conjecture.
- [7] Jannsen, U.: Motivic Sheaves and Filtrations on Chow Groups, in *Motives*, *Proc. Symp. Pure Math.*, Vol. **55** (1994), 245-302.
- [8] Kato, K., Saito, T.: Ramification theory for varieties over a perfect field, preprint 59 pages, 2004.
- [9] Kleinman, S: Algebraic Cycles and the Weil Conjectures, in *Dix Exposés sur la Cohomologie des Schémas*, North-Holland, Amsterdam, 1968, 359-386.
- [10] Kleinman, S: The Standard Conjectures, in *Motives*, *Proc. Symp. Pure Math.*, Vol. **55** (1994), 13-20.
- [11] Levine, M.: *Mixed Motives*, *Mathematical Surveys and Monographs* **57** (1998), American Mathematical Society.
- [12] Murre, J.P.: On a conjectural filtration on the Chow groups of an algebraic variety, (I and II), *Indag. Math. New Series* Vol. **4** (1993), 177-201.
- [13] Pink, R.: On the calculation of local terms in the Lefschetz-Verdier trace formula and its application to a conjecture of Deligne, *Ann. of Math.* **135** (1992), 483-525.
- [14] Zink, T.: The Lefschetz trace formula for open surfaces, In: *Automorphic Forms, Shimura Varieties and L -functions*, vol. II, *Prospectives in Math.* (1989), 337-376.

DEPT. OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, IL 60637,
USA

E-mail address: `bloch@math.uchicago.edu`

MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, FB6, MATHEMATIK, 45117
ESSEN, GERMANY

E-mail address: `esnault@uni-essen.de`