

SCHRIFTENREIHE DER FAKULTÄT FÜR MATHEMATIK

A Global div-curl-Lemma
for Mixed Boundary Conditions in Weak Lipschitz Domains
and a Corresponding Generalized A_0^* - A_1 -Lemma in Hilbert Spaces

by
Dirk Pauly

SM-UDE-812

2017

Received: July 17, 2017

A Global div-curl-Lemma for Mixed Boundary Conditions in Weak Lipschitz Domains and a Corresponding Generalized A_0^* - A_1 -Lemma in Hilbert Spaces

DIRK PAULY

ABSTRACT. We prove global and local versions of the so called div-curl-lemma, also known as compensated compactness, for mixed boundary conditions as well as bounded weak Lipschitz domains in 3D and weak Lipschitz interfaces. We will generalize our results using an abstract Hilbert space setting, which shows corresponding results to hold in arbitrary dimensions as well as for various differential operators. The crucial tools are Hilbert complexes and related compact embeddings.

CONTENTS

1.	Introduction	1
2.	Definitions and Preliminaries	2
3.	The div-rot-Lemma	4
4.	Generalizations	5
4.1.	Functional Analysis Toolbox	5
4.2.	The A_0^* - A_1 -Lemma	7
5.	Applications	8
5.1.	The div-rot-Lemma Revisited	8
5.2.	Generalized Electro-Magnetics	9
5.3.	Biharmonic Equation and General Relativity, Gravitational Waves	10
5.4.	Linear Elasticity	11
	References	12

1. INTRODUCTION

We will prove a global version of the so called div-curl-lemma used for compensated compactness, stating that under certain (mixed tangential and normal) boundary conditions and (very weak) regularity assumptions on a domain $\Omega \subset \mathbb{R}^3$ the following holds:

Let $\Omega \subset \mathbb{R}^3$ be a bounded weak Lipschitz domain with boundary Γ and boundary parts $\Gamma_{\mathbf{t}}$ and $\Gamma_{\mathbf{n}}$. Let (E_n) and (H_n) be two sequences bounded in $L^2(\Omega)$, such that $(\operatorname{curl} E_n)$ and $(\operatorname{div} H_n)$ are also bounded in $L^2(\Omega)$ and $\nu \times E_n = 0$ on $\Gamma_{\mathbf{t}}$ and $\nu \cdot H_n = 0$ on $\Gamma_{\mathbf{n}}$. Then there exist subsequences, again denoted by (E_n) and (H_n) , such that (E_n) , $(\operatorname{curl} E_n)$ and (H_n) , $(\operatorname{div} H_n)$ converge weakly to E , $\operatorname{curl} E$ resp. H , $\operatorname{div} H$ in $L^2(\Omega)$ and the inner products converge as well, i.e.,

$$\int_{\Omega} E_n \cdot H_n \rightarrow \int_{\Omega} E \cdot H.$$

A local version (distributional like convergence for arbitrary domains and no boundary conditions needed) of this div-curl-lemma is then immediately obtained.

Let $\Omega \subset \mathbb{R}^3$ be an open set. Let (E_n) and (H_n) be two sequences bounded in $L^2(\Omega)$, such that $(\operatorname{curl} E_n)$ and $(\operatorname{div} H_n)$ are also bounded in $L^2(\Omega)$. Then there exist subsequences, again denoted by (E_n) and (H_n) ,

Date: July 17, 2017.

1991 Mathematics Subject Classification. 35B27, 35Q61, 47B07, 46B50.

Key words and phrases. div-curl-lemma, compensated compactness, mixed boundary conditions, weak Lipschitz domains, Maxwell's equations.

such that (E_n) , $(\operatorname{curl} E_n)$ and (H_n) , $(\operatorname{div} H_n)$ converge weakly to E , $\operatorname{curl} E$ resp. H , $\operatorname{div} H$ in $L^2(\Omega)$ and the inner products converge in the distributional sense as well, i.e., for all $\varphi \in \dot{C}^\infty(\Omega)$ it holds

$$\int_{\Omega} (E_n \cdot H_n) \varphi \rightarrow \int_{\Omega} (E \cdot H) \varphi.$$

For details, see Theorem 3.1 and Corollary 3.2. We will also show a generalization to a natural Hilbert space setting in Theorem 4.7. In Section 5 we apply this result to some more differential operators in 3D and ND, appearing, e.g., in generalized electro-magnetics, for the biharmonic equation, in general relativity, for gravitational waves, and in the theory of linear elasticity and plasticity.

The div-curl-lemma, or compensated compactness, see the original papers by Murat [13] and Tartar [23] or [7, 22], and its variants and extensions have plenty of important applications. For an extensive discussion and a historical overview of the div-curl-lemma see [24]. More recent discussions can be found, e.g., in [5, 25] and in the nice preprint [26]. The div-curl-lemma is widely used in the theory of homogenization of (nonlinear) partial differential equations, see, e.g., [22]. Moreover, it is crucial in establishing compactness and regularity results for nonlinear partial differential equations such as harmonic maps, see, e.g., [9, 8, 19]. Numerical applications can be found, e.g., in [2]. It is further a crucial tool in the homogenization of stochastic partial differential equations, especially with certain random coefficients, see, e.g., the survey [1] and the literature cited therein, e.g., [10].

Let us also mention that the div-curl-lemma is particularly useful to treat homogenization of problems arising in plasticity, see, e.g., a recent preprint on this topic [21], for which the preprint [20] provides the important key div-curl-lemma. As in [20, 21] $H^1(\Omega)$ -potentials are used, these contributions are restricted to smooth, e.g., C^2 or convex, domains and to full boundary conditions. On the other hand, using Weck's selection theorem (2.1) it is easily possible to extend these results even to bounded weak Lipschitz domains of arbitrary topology and to the case of mixed boundary conditions.

Generally, for problems related to Maxwell's equations the detour over $H^1(\Omega)$ instead of using Weck's selection theorem seems to be the wrong way to deal with such equations. Most of the arguments simply fail, and if not, the results are usually limited to smooth domains and trivial topologies. Mixed boundary conditions cannot be treated properly. Since the early 1970's, see the original paper by Weck [28] for Weck's selection theorem, it is well known, that the $H^1(\Omega)$ -detour is often not helpful and does not lead to satisfying results.

2. DEFINITIONS AND PRELIMINARIES

Let $\Omega \subset \mathbb{R}^3$ be a bounded weak Lipschitz domain, see [3, Definition 2.3] for details, with boundary $\Gamma := \partial\Omega$, which is divided into two relatively open weak Lipschitz subsets Γ_t and $\Gamma_n := \Gamma \setminus \overline{\Gamma_t}$ (its complement), see [3, Definition 2.5] for details. Note that strong Lipschitz (graph of Lipschitz functions) implies weak Lipschitz (Lipschitz manifolds) for the boundary as well as the interface. Throughout this paper we shall assume the latter regularity on Ω and Γ_t .

Recently, in [3], Weck's selection theorem, also known as the Maxwell compactness property, has been shown to hold for such bounded weak Lipschitz domains and mixed boundary conditions. More precisely, the embedding

$$(2.1) \quad \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega)$$

is compact, see [3, Theorem 4.7]. A short historical overview of Weck's selection theorem is given in the introduction of [3], see also the original paper [28] and [18, 27, 6, 29, 11, 12] for simpler proofs and generalizations.

Here the usual Lebesgue and Sobolev spaces are denoted by $L^2(\Omega)$ and $H^1(\Omega)$ as well as

$$R(\Omega) := \{E \in L^2(\Omega) : \operatorname{rot} E \in L^2(\Omega)\}, \quad D(\Omega) := \{E \in L^2(\Omega) : \operatorname{div} E \in L^2(\Omega)\},$$

where we prefer to write rot instead of curl . $R(\Omega)$ and $D(\Omega)$ are also written as $H(\operatorname{rot}, \Omega)$ or $H(\operatorname{curl}, \Omega)$ resp. $H(\operatorname{div}, \Omega)$ in the literature. With the help of test functions and test vector fields

$$\mathring{C}_{\Gamma_t}^\infty(\Omega) := \{\varphi|_{\Omega} : \varphi \in \dot{C}^\infty(\mathbb{R}^3), \operatorname{dist}(\operatorname{supp} \varphi, \Gamma_t) > 0\}$$

we define the closed subspaces

$$(2.2) \quad \mathring{H}_{\Gamma_t}^1(\Omega) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega)}^{\mathbf{H}^1(\Omega)}, \quad \mathring{R}_{\Gamma_t}(\Omega) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega)}^{\mathbf{R}(\Omega)}, \quad \mathring{D}_{\Gamma_n}(\Omega) := \overline{\mathring{C}_{\Gamma_n}^\infty(\Omega)}^{\mathbf{D}(\Omega)}$$

as closures of test functions respectively vector fields. In (2.2) homogeneous scalar, tangential and normal traces on Γ_t resp. Γ_n are generalized, respectively. To avoid case studies when using the Poincaré estimate, we also define

$$\mathring{H}_\emptyset^1(\Omega) := \mathbf{H}^1(\Omega) \cap \mathbb{R}^{\perp L^2(\Omega)} = \left\{ u \in \mathbf{H}^1(\Omega) : \int_\Omega u = 0 \right\}.$$

Let us emphasize that our assumptions also allow for Rellich's selection theorem, i.e., the embedding

$$(2.3) \quad \mathring{H}_{\Gamma_t}^1(\Omega) \hookrightarrow \mathbf{L}^2(\Omega)$$

is compact, see, e.g., [3, Theorem 4.8]. By density we have the two rules of partial integration

$$(2.4) \quad \forall u \in \mathring{H}_{\Gamma_t}^1(\Omega) \quad \forall H \in \mathring{D}_{\Gamma_n}(\Omega) \quad \langle \nabla u, H \rangle_{\mathbf{L}^2(\Omega)} = -\langle u, \operatorname{div} H \rangle_{\mathbf{L}^2(\Omega)},$$

$$(2.5) \quad \forall E \in \mathring{R}_{\Gamma_t}(\Omega) \quad \forall H \in \mathring{R}_{\Gamma_n}(\Omega) \quad \langle \operatorname{rot} E, H \rangle_{\mathbf{L}^2(\Omega)} = \langle E, \operatorname{rot} H \rangle_{\mathbf{L}^2(\Omega)}.$$

We emphasize that, besides Weck's selection theorem, the resulting Maxwell estimates (Friedrichs/Poincaré type estimates), Helmholtz decompositions, closed ranges, continuous and compact inverse operators, and an adequate electro-magneto static solution theory for bounded weak Lipschitz domains and mixed boundary conditions, another important result has been shown in [3]. It holds

$$(2.6) \quad \begin{aligned} \mathring{H}_{\Gamma_t}^1(\Omega) &= \left\{ u \in \mathbf{H}^1(\Omega) : \langle \nabla u, \Phi \rangle_{\mathbf{L}^2(\Omega)} = -\langle u, \operatorname{div} \Phi \rangle_{\mathbf{L}^2(\Omega)} \text{ for all } \Phi \in \mathring{C}_{\Gamma_n}^\infty(\Omega) \right\}, \\ \mathring{R}_{\Gamma_t}(\Omega) &= \left\{ E \in \mathbf{R}(\Omega) : \langle \operatorname{rot} E, \Phi \rangle_{\mathbf{L}^2(\Omega)} = \langle E, \operatorname{rot} \Phi \rangle_{\mathbf{L}^2(\Omega)} \text{ for all } \Phi \in \mathring{C}_{\Gamma_n}^\infty(\Omega) \right\}, \\ \mathring{D}_{\Gamma_n}(\Omega) &= \left\{ H \in \mathbf{D}(\Omega) : \langle \operatorname{div} H, \varphi \rangle_{\mathbf{L}^2(\Omega)} = -\langle H, \nabla \varphi \rangle_{\mathbf{L}^2(\Omega)} \text{ for all } \varphi \in \mathring{C}_{\Gamma_t}^\infty(\Omega) \right\}, \end{aligned}$$

i.e., strong and weak definitions of boundary conditions coincide, see [3, Theorem 4.5]. Furthermore, we define the closed subspaces of irrotational resp. solenoidal vector fields

$$\mathbf{R}_0(\Omega) := \{ E \in \mathbf{R}(\Omega) : \operatorname{rot} E = 0 \}, \quad \mathbf{D}_0(\Omega) := \{ E \in \mathbf{D}(\Omega) : \operatorname{div} E = 0 \}$$

as well as

$$\mathring{R}_{\Gamma_t,0}(\Omega) := \mathring{R}_{\Gamma_t}(\Omega) \cap \mathbf{R}_0(\Omega), \quad \mathring{D}_{\Gamma_n,0}(\Omega) := \mathring{D}_{\Gamma_n}(\Omega) \cap \mathbf{D}_0(\Omega).$$

A direct consequence of (2.1) is the compactness of the unit ball in

$$\mathcal{H}(\Omega) := \mathring{R}_{\Gamma_t,0}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega),$$

the space of so called Dirichlet-Neumann fields. Hence $\mathcal{H}(\Omega)$ is finite dimensional. Another immediate consequence of Weck's selection theorem (2.1), using a standard indirect argument, is the so called Maxwell estimate, i.e.,

$$(2.7) \quad \exists c_m > 0 \quad \forall E \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n}(\Omega) \cap \mathcal{H}(\Omega)^{\perp L^2(\Omega)} \quad |E|_{\mathbf{L}^2(\Omega)} \leq c_m (|\operatorname{rot} E|_{\mathbf{L}^2(\Omega)} + |\operatorname{div} E|_{\mathbf{L}^2(\Omega)})$$

or, equivalently,

$$(2.8) \quad \forall E \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n}(\Omega) \quad |E - \pi E|_{\mathbf{L}^2(\Omega)} \leq c_m (|\operatorname{rot} E|_{\mathbf{L}^2(\Omega)} + |\operatorname{div} E|_{\mathbf{L}^2(\Omega)}),$$

see [3, Theorem 5.1], where $\pi : \mathbf{L}^2(\Omega) \rightarrow \mathcal{H}(\Omega)$ denotes the $\mathbf{L}^2(\Omega)$ -orthonormal projector onto the Dirichlet-Neumann fields. Recent estimates for the Maxwell constant c_m can be found in [14, 15, 16]. Analogously, Rellich's selection theorem (2.3) shows the Friedrichs/Poincaré estimate

$$(2.9) \quad \exists c_{f,p} > 0 \quad \forall u \in \mathring{H}_{\Gamma_t}^1(\Omega) \quad |u|_{\mathbf{L}^2(\Omega)} \leq c_{f,p} |\nabla u|_{\mathbf{L}^2(\Omega)},$$

see [3, Theorem 4.8]. By the projection theorem, applied to the densely defined and closed (unbounded) linear operator

$$\nabla : \mathring{H}_{\Gamma_t}^1(\Omega) \subset \mathbf{L}^2(\Omega) \longrightarrow \mathbf{L}^2(\Omega)$$

with (Hilbert space) adjoint

$$\nabla^* = -\operatorname{div} : \mathring{D}_{\Gamma_n}(\Omega) \subset L^2(\Omega) \longrightarrow L^2(\Omega),$$

here we need (2.6), we get the simple Helmholtz decomposition

$$(2.10) \quad L^2(\Omega) = \nabla \mathring{H}_{\Gamma_t}^1(\Omega) \oplus_{L^2(\Omega)} \mathring{D}_{\Gamma_n,0}(\Omega),$$

see [3, Theorem 5.3 or (13)], which immediately implies

$$(2.11) \quad \mathring{R}_{\Gamma_t}(\Omega) = \nabla \mathring{H}_{\Gamma_t}^1(\Omega) \oplus_{L^2(\Omega)} (\mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega))$$

as $\nabla \mathring{H}_{\Gamma_t}^1(\Omega) \subset \mathring{R}_{\Gamma_t,0}(\Omega)$. Note that the decompositions (2.10) and (2.11) are orthogonal which is denoted by $\oplus_{L^2(\Omega)}$. By (2.9), the range $\nabla \mathring{H}_{\Gamma_t}^1(\Omega)$ is closed in $L^2(\Omega)$, see also [3, Lemma 5.2]. Note that we call (2.10) a simple Helmholtz decomposition, since the refined Helmholtz decomposition

$$L^2(\Omega) = \nabla \mathring{H}_{\Gamma_t}^1(\Omega) \oplus_{L^2(\Omega)} \mathcal{H}(\Omega) \oplus_{L^2(\Omega)} \operatorname{rot} \mathring{R}_{\Gamma_n}(\Omega)$$

holds as well, see [3, Theorem 5.3], where also $\operatorname{rot} \mathring{R}_{\Gamma_n}(\Omega)$ is closed in $L^2(\Omega)$ as a consequence of (2.7), see [3, Lemma 5.2].

3. THE DIV-ROT-LEMMA

Theorem 3.1 (global div-rot-lemma). *Let $(E_n) \subset \mathring{R}_{\Gamma_t}(\Omega)$ and $(H_n) \subset \mathring{D}_{\Gamma_n}(\Omega)$ be two sequences bounded in $R(\Omega)$ resp. $D(\Omega)$. Then there exist $E \in \mathring{R}_{\Gamma_t}(\Omega)$ and $H \in \mathring{D}_{\Gamma_n}(\Omega)$ as well as subsequences, again denoted by (E_n) and (H_n) , such that (E_n) and (H_n) converge weakly in $R(\Omega)$ resp. $D(\Omega)$ to E resp. H together with the convergence of the inner products*

$$\langle E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle E, H \rangle_{L^2(\Omega)}.$$

Proof. We pick subsequences, again denoted by (E_n) and (H_n) , such that (E_n) and (H_n) converge weakly in $R(\Omega)$ resp. $D(\Omega)$ to E resp. H for some $E \in \mathring{R}_{\Gamma_t}(\Omega)$ and $H \in \mathring{D}_{\Gamma_n}(\Omega)$. By the simple Helmholtz decomposition (2.11), we have the orthogonal decomposition $\mathring{R}_{\Gamma_t}(\Omega) \ni E_n = \nabla u_n + \tilde{E}_n$ with some $u_n \in \mathring{H}_{\Gamma_t}^1(\Omega)$ and $\tilde{E}_n \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega)$. Then (u_n) is bounded in $H^1(\Omega)$ by orthogonality and the Friedrichs/Poincaré estimate (2.9). (\tilde{E}_n) is bounded in $R(\Omega) \cap D(\Omega)$ by orthogonality and $\operatorname{rot} \tilde{E}_n = \operatorname{rot} E_n$, $\operatorname{div} \tilde{E}_n = 0$. Hence, using Rellich's and Weck's selection theorems there exist $u \in \mathring{H}_{\Gamma_t}^1(\Omega)$ and $\tilde{E} \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega)$ and we can extract two subsequences, again denoted by (u_n) and (\tilde{E}_n) , such that $u_n \rightharpoonup u$ in $H^1(\Omega)$ and $u_n \rightarrow u$ in $L^2(\Omega)$ as well as $\tilde{E}_n \rightharpoonup \tilde{E}$ in $R(\Omega) \cap D(\Omega)$ and $\tilde{E}_n \rightarrow \tilde{E}$ in $L^2(\Omega)$. We have $E = \nabla u + \tilde{E}$, giving the simple Helmholtz decomposition for E , as, e.g., for all $\varphi \in \mathring{C}^\infty(\Omega)$

$$\langle E, \varphi \rangle_{L^2(\Omega)} \leftarrow \langle E_n, \varphi \rangle_{L^2(\Omega)} = \langle \nabla u_n, \varphi \rangle_{L^2(\Omega)} + \langle \tilde{E}_n, \varphi \rangle_{L^2(\Omega)} \rightarrow \langle \nabla u, \varphi \rangle_{L^2(\Omega)} + \langle \tilde{E}, \varphi \rangle_{L^2(\Omega)}.$$

Then by (2.4)

$$\begin{aligned} \langle E_n, H_n \rangle_{L^2(\Omega)} &= \langle \nabla u_n, H_n \rangle_{L^2(\Omega)} + \langle \tilde{E}_n, H_n \rangle_{L^2(\Omega)} = -\langle u_n, \operatorname{div} H_n \rangle_{L^2(\Omega)} + \langle \tilde{E}_n, H_n \rangle_{L^2(\Omega)} \\ &\rightarrow -\langle u, \operatorname{div} H \rangle_{L^2(\Omega)} + \langle \tilde{E}, H \rangle_{L^2(\Omega)} = \langle \nabla u, H \rangle_{L^2(\Omega)} + \langle \tilde{E}, H \rangle_{L^2(\Omega)} = \langle E, H \rangle_{L^2(\Omega)}, \end{aligned}$$

completing the proof. \square

Corollary 3.2 (local div-rot-lemma). *Let $(E_n) \subset R(\Omega)$ and $(H_n) \subset D(\Omega)$ be two sequences bounded in $R(\Omega)$ resp. $D(\Omega)$. Then there exist $E \in R(\Omega)$ and $H \in D(\Omega)$ as well as subsequences, again denoted by (E_n) and (H_n) , such that $E_n \rightharpoonup E$ in $R(\Omega)$ and $H_n \rightharpoonup H$ in $D(\Omega)$ together with the distributional convergence*

$$\forall \varphi \in \mathring{C}^\infty(\Omega) \quad \langle \varphi E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle \varphi E, H \rangle_{L^2(\Omega)}.$$

Proof. Let $\Gamma_t := \Gamma$ and hence $\Gamma_n = \emptyset$. (φE_n) is bounded in $\mathring{R}_\Gamma(\Omega)$ and (H_n) is bounded in $D(\Omega)$. Theorem 3.1 shows the assertion. \square

Remark 3.3. We note that the boundedness of (E_n) and (H_n) in local spaces is sufficient for Corollary 3.2 to hold. Hence, no regularity or boundedness assumptions on Ω are needed, i.e., Ω can be an arbitrary domain in \mathbb{R}^3 .

4. GENERALIZATIONS

The idea of the proof of Theorem 3.1 can be generalized.

4.1. Functional Analysis Toolbox. Let $A : D(A) \subset H_1 \rightarrow H_2$ be a (possibly unbounded) closed and densely defined linear operator on two Hilbert spaces H_1 and H_2 with adjoint $A^* : D(A^*) \subset H_2 \rightarrow H_1$. Note $(A^*)^* = \overline{A} = A$, i.e., (A, A^*) is a dual pair. By the projection theorem the Helmholtz type decompositions

$$(4.1) \quad H_1 = N(A) \oplus_{H_1} \overline{R(A^*)}, \quad H_2 = N(A^*) \oplus_{H_2} \overline{R(A)}$$

hold, where we introduce the notation N for the kernel (or null space) and R for the range of a linear operator. We can define the reduced operators

$$\begin{aligned} \mathcal{A} &:= A|_{\overline{R(A^*)}} : D(\mathcal{A}) \subset \overline{R(A^*)} \rightarrow \overline{R(A)}, & D(\mathcal{A}) &:= D(A) \cap N(A)^{\perp_{H_1}} = D(A) \cap \overline{R(A^*)}, \\ \mathcal{A}^* &:= A^*|_{\overline{R(A)}} : D(\mathcal{A}^*) \subset \overline{R(A)} \rightarrow \overline{R(A^*)}, & D(\mathcal{A}^*) &:= D(A^*) \cap N(A^*)^{\perp_{H_2}} = D(A^*) \cap \overline{R(A)}, \end{aligned}$$

which are also closed and densely defined linear operators. We note that \mathcal{A} and \mathcal{A}^* are indeed adjoint to each other, i.e., $(\mathcal{A}, \mathcal{A}^*)$ is a dual pair as well. Now the inverse operators

$$\mathcal{A}^{-1} : R(\mathcal{A}) \rightarrow D(\mathcal{A}), \quad (\mathcal{A}^*)^{-1} : R(\mathcal{A}^*) \rightarrow D(\mathcal{A}^*)$$

exist and they are bijective, since \mathcal{A} and \mathcal{A}^* are injective by definition. Furthermore, by (4.1) we have the refined Helmholtz type decompositions

$$(4.2) \quad D(A) = N(A) \oplus_{H_1} D(\mathcal{A}), \quad D(A^*) = N(A^*) \oplus_{H_2} D(\mathcal{A}^*)$$

and thus we obtain for the ranges

$$(4.3) \quad R(A) = R(\mathcal{A}), \quad R(A^*) = R(\mathcal{A}^*).$$

By the closed range theorem and the closed graph theorem we get immediately the following.

Lemma 4.1. *The following assertions are equivalent:*

- (i) $\exists c_A \in (0, \infty) \quad \forall x \in D(\mathcal{A}) \quad |x|_{H_1} \leq c_A |Ax|_{H_2}$
- (i*) $\exists c_{A^*} \in (0, \infty) \quad \forall y \in D(\mathcal{A}^*) \quad |y|_{H_2} \leq c_{A^*} |A^*y|_{H_1}$
- (ii) $R(A) = R(\mathcal{A})$ is closed in H_2 .
- (ii*) $R(A^*) = R(\mathcal{A}^*)$ is closed in H_1 .
- (iii) $\mathcal{A}^{-1} : R(\mathcal{A}) \rightarrow D(\mathcal{A})$ is continuous and bijective with norm bounded by $(1 + c_A^2)^{1/2}$.
- (iii*) $(\mathcal{A}^*)^{-1} : R(\mathcal{A}^*) \rightarrow D(\mathcal{A}^*)$ is continuous and bijective with norm bounded by $(1 + c_{A^*}^2)^{1/2}$.

In case that one of the latter assertions is true, e.g., (ii), $R(A)$ is closed, we have

$$\begin{aligned} H_1 &= N(A) \oplus_{H_1} R(A^*), & H_2 &= N(A^*) \oplus_{H_2} R(A), \\ D(A) &= N(A) \oplus_{H_1} D(\mathcal{A}), & D(A^*) &= N(A^*) \oplus_{H_2} D(\mathcal{A}^*), \\ D(\mathcal{A}) &= D(A) \cap R(A^*), & D(\mathcal{A}^*) &= D(A^*) \cap R(A), \end{aligned}$$

and

$$A : D(\mathcal{A}) \subset R(A^*) \rightarrow R(A), \quad A^* : D(\mathcal{A}^*) \subset R(A) \rightarrow R(A^*).$$

Remark 4.2. For the “best” constants c_A, c_{A^*} the following holds: The Rayleigh quotients

$$\frac{1}{c_A} := \inf_{0 \neq x \in D(\mathcal{A})} \frac{|Ax|_{H_2}}{|x|_{H_1}}, \quad \frac{1}{c_{A^*}} := \inf_{0 \neq y \in D(\mathcal{A}^*)} \frac{|A^*y|_{H_1}}{|y|_{H_2}}$$

coincide, i.e., $c_A = c_{A^*} \in (0, \infty]$.

Lemma 4.3. *The following assertions are equivalent:*

- (i) $D(\mathcal{A}) \overset{c}{\rightleftarrows} \mathbf{H}_1$ is compact.
- (i*) $D(\mathcal{A}^*) \overset{c}{\rightleftarrows} \mathbf{H}_2$ is compact.
- (ii) $\mathcal{A}^{-1} : R(\mathbf{A}) \rightarrow R(\mathbf{A}^*)$ is compact.
- (ii*) $(\mathcal{A}^*)^{-1} : R(\mathbf{A}^*) \rightarrow R(\mathbf{A})$ is compact.

If one of these assertions holds true, e.g., (i), $D(\mathcal{A}) \overset{c}{\rightleftarrows} \mathbf{H}_1$ is compact, then the assertions of Lemma 4.1 and Remark 4.2 hold with $c_{\mathbf{A}} = c_{\mathbf{A}^*} \in (0, \infty)$. Especially, the Friedrichs/Poincaré type estimates hold, all ranges are closed and the inverse operators

$$\mathcal{A}^{-1} : R(\mathbf{A}) \rightarrow R(\mathbf{A}^*), \quad (\mathcal{A}^*)^{-1} : R(\mathbf{A}^*) \rightarrow R(\mathbf{A})$$

are compact with norms $\|\mathcal{A}^{-1}\|_{R(\mathbf{A}), R(\mathbf{A}^*)} = \|(\mathcal{A}^*)^{-1}\|_{R(\mathbf{A}^*), R(\mathbf{A})} = c_{\mathbf{A}}$.

Proof. As the other assertions are easily proved or immediately clear by symmetry, we just show that (i), i.e., the compactness of

$$D(\mathcal{A}) = D(\mathbf{A}) \cap \overline{R(\mathbf{A}^*)} \overset{c}{\rightleftarrows} \mathbf{H}_1,$$

implies (i*) as well as Lemma 4.1 (i).

(i) \Rightarrow Lemma 4.1 (i): For this we use a standard indirect argument. If Lemma 4.1 (i) was wrong, there exists a sequence $(x_n) \subset D(\mathcal{A})$ with $\|x_n\|_{\mathbf{H}_1} = 1$ and $\mathbf{A}x_n \rightarrow 0$. As (x_n) is bounded in $D(\mathcal{A})$ we can extract a subsequence, again denoted by (x_n) , with $x_n \rightarrow x \in \mathbf{H}_1$ in \mathbf{H}_1 . Since \mathcal{A} is closed, we have $x \in D(\mathcal{A})$ and $\mathbf{A}x = 0$, hence $x \in N(\mathcal{A}) = \{0\}$, in contradiction to $1 = \|x_n\|_{\mathbf{H}_1} \rightarrow \|x\|_{\mathbf{H}_1} = 0$.

(i) \Rightarrow (i*): Let $(y_n) \subset D(\mathcal{A}^*)$ be a bounded sequence. Utilizing Lemma 4.1 (i) and (ii) we obtain $D(\mathcal{A}^*) = D(\mathbf{A}^*) \cap R(\mathbf{A})$ and thus $y_n = \mathbf{A}x_n$ with $(x_n) \subset D(\mathcal{A})$, which is bounded in $D(\mathcal{A})$ by Lemma 4.1 (i). Hence we may extract a subsequence, again denoted by (x_n) , converging in \mathbf{H}_1 . Therefore with $x_{n,m} := x_n - x_m$ and $y_{n,m} := y_n - y_m$ we see

$$\|y_{n,m}\|_{\mathbf{H}_2}^2 = \langle y_{n,m}, \mathbf{A}(x_{n,m}) \rangle_{\mathbf{H}_2} = \langle \mathbf{A}^*(y_{n,m}), x_{n,m} \rangle_{\mathbf{H}_1} \leq c \|x_{n,m}\|_{\mathbf{H}_1},$$

and hence (y_n) is a Cauchy sequence in \mathbf{H}_2 . \square

Now, let $\mathbf{A}_0 : D(\mathbf{A}_0) \subset \mathbf{H}_0 \rightarrow \mathbf{H}_1$ and $\mathbf{A}_1 : D(\mathbf{A}_1) \subset \mathbf{H}_1 \rightarrow \mathbf{H}_2$ be (possibly unbounded) closed and densely defined linear operators on three Hilbert spaces \mathbf{H}_0 , \mathbf{H}_1 , and \mathbf{H}_2 with adjoints $\mathbf{A}_0^* : D(\mathbf{A}_0^*) \subset \mathbf{H}_1 \rightarrow \mathbf{H}_0$ and $\mathbf{A}_1^* : D(\mathbf{A}_1^*) \subset \mathbf{H}_2 \rightarrow \mathbf{H}_1$ as well as reduced operators \mathcal{A}_0 , \mathcal{A}_0^* , and \mathcal{A}_1 , \mathcal{A}_1^* . Furthermore, we assume the sequence or complex property of \mathbf{A}_0 and \mathbf{A}_1 , that is, $\mathbf{A}_1 \mathbf{A}_0 = 0$, i.e.,

$$(4.4) \quad R(\mathbf{A}_0) \subset N(\mathbf{A}_1).$$

Then also $\mathbf{A}_0^* \mathbf{A}_1^* = 0$, i.e., $R(\mathbf{A}_1^*) \subset N(\mathbf{A}_0^*)$. From the Helmholtz type decompositions (4.1) for $\mathbf{A} = \mathbf{A}_0$ and $\mathbf{A} = \mathbf{A}_1$ we get in particular

$$(4.5) \quad \mathbf{H}_1 = \overline{R(\mathbf{A}_0)} \oplus_{\mathbf{H}_1} N(\mathbf{A}_0^*), \quad \mathbf{H}_1 = \overline{R(\mathbf{A}_1^*)} \oplus_{\mathbf{H}_1} N(\mathbf{A}_1),$$

and the following result for Helmholtz type decompositions:

Lemma 4.4. *Let $N_{0,1} := N(\mathbf{A}_1) \cap N(\mathbf{A}_0^*)$. The refined Helmholtz type decompositions*

$$(4.6) \quad N(\mathbf{A}_1) = \overline{R(\mathbf{A}_0)} \oplus_{\mathbf{H}_1} N_{0,1}, \quad D(\mathbf{A}_1) = \overline{R(\mathbf{A}_0)} \oplus_{\mathbf{H}_1} (D(\mathbf{A}_1) \cap N(\mathbf{A}_0^*)), \quad R(\mathbf{A}_0) = R(\mathcal{A}_0),$$

$$(4.7) \quad N(\mathbf{A}_0^*) = \overline{R(\mathbf{A}_1^*)} \oplus_{\mathbf{H}_1} N_{0,1}, \quad D(\mathbf{A}_0^*) = \overline{R(\mathbf{A}_1^*)} \oplus_{\mathbf{H}_1} (D(\mathbf{A}_0^*) \cap N(\mathbf{A}_1)), \quad R(\mathbf{A}_1^*) = R(\mathcal{A}_1^*),$$

and

$$(4.8) \quad \mathbf{H}_1 = \overline{R(\mathbf{A}_0)} \oplus_{\mathbf{H}_1} N_{0,1} \oplus_{\mathbf{H}_1} \overline{R(\mathbf{A}_1^*)}$$

hold, which can be further refined and specialized, e.g., to

$$(4.9) \quad D(\mathbf{A}_1) \cap D(\mathbf{A}_0^*) = D(\mathcal{A}_0^*) \oplus_{\mathbf{H}_1} N_{0,1} \oplus_{\mathbf{H}_1} D(\mathcal{A}_1).$$

Proof. By (4.5) and the complex properties we see (4.6) and (4.7), yielding directly (4.8) and (4.9). \square

We observe

$$\begin{aligned} D(\mathcal{A}_1) &= D(\mathcal{A}_1) \cap \overline{R(\mathcal{A}_1^*)} \subset D(\mathcal{A}_1) \cap N(\mathcal{A}_0^*) \subset D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*), \\ D(\mathcal{A}_0^*) &= D(\mathcal{A}_0^*) \cap \overline{R(\mathcal{A}_0)} \subset D(\mathcal{A}_0^*) \cap N(\mathcal{A}_1) \subset D(\mathcal{A}_0^*) \cap D(\mathcal{A}_1), \end{aligned}$$

and using the refined Helmholtz type decompositions of Lemma 4.4 as well as the results of Lemma 4.1, Lemma 4.3, and Lemma 4.5, we immediately see:

Lemma 4.5. *The following assertions are equivalent:*

- (i) $D(\mathcal{A}_0) \overset{\circ}{\iff} \mathbf{H}_0$, $D(\mathcal{A}_1) \overset{\circ}{\iff} \mathbf{H}_1$, and $N_{0,1} \overset{\circ}{\iff} \mathbf{H}_1$ are compact.
- (ii) $D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*) \overset{\circ}{\iff} \mathbf{H}_1$ is compact.

In this case, the cohomology group $N_{0,1}$ has finite dimension.

We summarize:

Theorem 4.6. *Let $D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*) \overset{\circ}{\iff} \mathbf{H}_1$ be compact. Then $D(\mathcal{A}_0) \overset{\circ}{\iff} \mathbf{H}_0$, $D(\mathcal{A}_1) \overset{\circ}{\iff} \mathbf{H}_1$, and $D(\mathcal{A}_0^*) \overset{\circ}{\iff} \mathbf{H}_1$, $D(\mathcal{A}_1^*) \overset{\circ}{\iff} \mathbf{H}_2$ are compact, all ranges $R(\mathcal{A}_0)$, $R(\mathcal{A}_0^*)$, and $R(\mathcal{A}_1)$, $R(\mathcal{A}_1^*)$ are closed, and the corresponding Friedrichs/Poincaré type estimates hold, i.e. there exists $c_{\mathcal{A}_0}, c_{\mathcal{A}_1} \in (0, \infty)$ such that*

$$(4.10) \quad \begin{aligned} \forall z \in D(\mathcal{A}_0) & \quad |z|_{\mathbf{H}_0} \leq c_{\mathcal{A}_0} |A_0 z|_{\mathbf{H}_1}, \\ \forall x \in D(\mathcal{A}_0^*) & \quad |x|_{\mathbf{H}_1} \leq c_{\mathcal{A}_0} |A_0^* x|_{\mathbf{H}_0}, \\ \forall x \in D(\mathcal{A}_1) & \quad |x|_{\mathbf{H}_1} \leq c_{\mathcal{A}_1} |A_1 x|_{\mathbf{H}_2}, \\ \forall y \in D(\mathcal{A}_1^*) & \quad |y|_{\mathbf{H}_2} \leq c_{\mathcal{A}_1} |A_1^* y|_{\mathbf{H}_1}. \end{aligned}$$

Moreover, all refined Helmholtz type decompositions of Lemma 4.4 hold with closed ranges, especially

$$(4.11) \quad D(\mathcal{A}_1) = R(\mathcal{A}_0) \oplus_{\mathbf{H}_1} (D(\mathcal{A}_1) \cap N(\mathcal{A}_0^*)).$$

Proof. Apply the latter lemmas and remarks to $\mathbf{A} = \mathbf{A}_0$ and $\mathbf{A} = \mathbf{A}_1$. □

4.2. The \mathbf{A}_0^* - \mathbf{A}_1 -Lemma. Let \mathbf{A}_0 and \mathbf{A}_1 be as introduced before satisfying the complex property (4.4), i.e., $\mathbf{A}_1 \mathbf{A}_0 = 0$ or $R(\mathbf{A}_0) \subset N(\mathbf{A}_1)$. In other words, the primal and dual sequences

$$(4.12) \quad \begin{aligned} D(\mathbf{A}_0) \subset \mathbf{H}_0 & \xrightarrow{A_0} D(\mathbf{A}_1) \subset \mathbf{H}_1 \xrightarrow{A_1} \mathbf{H}_2, \\ \mathbf{H}_0 \xleftarrow{A_0^*} D(\mathbf{A}_0^*) \subset \mathbf{H}_1 & \xleftarrow{A_1^*} D(\mathbf{A}_1^*) \subset \mathbf{H}_2 \end{aligned}$$

are Hilbert complexes of closed and densely defined linear operators. The additional assumption of closed ranges $R(\mathbf{A}_0)$ and $R(\mathbf{A}_1)$ (and hence also closed ranges $R(\mathbf{A}_0^*)$ and $R(\mathbf{A}_1^*)$) is equivalent to calling the Hilbert complexes closed. The complexes are exact, if and only if $N_{0,1} = \{0\}$.

As our main result, the following generalized global div-curl-lemma holds.

Theorem 4.7 (\mathbf{A}_0^* - \mathbf{A}_1 -lemma). *Let $D(\mathbf{A}_1) \cap D(\mathbf{A}_0^*) \overset{\circ}{\iff} \mathbf{H}_1$ be compact. Moreover, let $(x_n) \subset D(\mathbf{A}_1)$ and $(y_n) \subset D(\mathbf{A}_0^*)$ be two sequences bounded in $D(\mathbf{A}_1)$ resp. $D(\mathbf{A}_0^*)$. Then there exist $x \in D(\mathbf{A}_1)$ and $y \in D(\mathbf{A}_0^*)$ as well as subsequences, again denoted by (x_n) and (y_n) , such that (x_n) and (y_n) converge weakly in $D(\mathbf{A}_1)$ resp. $D(\mathbf{A}_0^*)$ to x resp. y together with the convergence of the inner products*

$$\langle x_n, y_n \rangle_{\mathbf{H}_1} \rightarrow \langle x, y \rangle_{\mathbf{H}_1}.$$

Proof. Note that Theorem 4.6 can be applied. We pick subsequences, again denoted by (x_n) and (y_n) , such that (x_n) and (y_n) converge weakly in $D(\mathbf{A}_1)$ resp. $D(\mathbf{A}_0^*)$ to $x \in D(\mathbf{A}_1)$ resp. $y \in D(\mathbf{A}_0^*)$. By (4.11) we get the orthogonal decomposition

$$D(\mathbf{A}_1) \ni x_n = A_0 z_n + \tilde{x}_n, \quad z_n \in D(\mathcal{A}_0), \quad \tilde{x}_n \in D(\mathbf{A}_1) \cap N(\mathbf{A}_0^*).$$

(z_n) is bounded in $D(\mathcal{A}_0)$ by orthogonality and the Friedrichs/Poincaré type estimate (4.10). (\tilde{x}_n) is bounded in $D(\mathbf{A}_1) \cap D(\mathbf{A}_0^*)$ by orthogonality and $A_1 \tilde{x}_n = A_1 x_n$, $A_0^* \tilde{x}_n = 0$. Using the compact embeddings $D(\mathcal{A}_0) \overset{\circ}{\iff} \mathbf{H}_0$ and $D(\mathbf{A}_1) \cap D(\mathbf{A}_0^*) \overset{\circ}{\iff} \mathbf{H}_1$, there exist $z \in D(\mathcal{A}_0)$ and $\tilde{x} \in D(\mathbf{A}_1) \cap N(\mathbf{A}_0^*)$ and we can extract two subsequences, again denoted by (z_n) and (\tilde{x}_n) , such that $z_n \rightarrow z$ in $D(\mathcal{A}_0)$ and $z_n \rightarrow z$ in

H_0 as well as $\tilde{x}_n \rightarrow \tilde{x}$ in $D(A_1) \cap D(A_0^*)$ and $\tilde{x}_n \rightarrow \tilde{x}$ in H_1 . We have $x = A_0 z + \tilde{x}$, giving the Helmholtz type decomposition for x , as, e.g., for all $\varphi \in H_1$

$$\langle x, \varphi \rangle_{H_1} \leftarrow \langle x_n, \varphi \rangle_{H_1} = \langle A_0 z_n, \varphi \rangle_{H_1} + \langle \tilde{x}_n, \varphi \rangle_{H_1} \rightarrow \langle A_0 z, \varphi \rangle_{H_1} + \langle \tilde{x}, \varphi \rangle_{H_1}.$$

Finally, we see

$$\begin{aligned} \langle x_n, y_n \rangle_{H_1} &= \langle A_0 z_n, y_n \rangle_{H_1} + \langle \tilde{x}_n, y_n \rangle_{H_1} = \langle z_n, A_0^* y_n \rangle_{H_0} + \langle \tilde{x}_n, y_n \rangle_{H_1} \\ &\rightarrow \langle z, A_0^* y \rangle_{H_0} + \langle \tilde{x}, y \rangle_{H_1} = \langle A_0 z, y \rangle_{H_1} + \langle \tilde{x}, y \rangle_{H_1} = \langle x, y \rangle_{H_1}, \end{aligned}$$

completing the proof. \square

Remark 4.8. *By Lemma 4.5 the crucial assumption, i.e., $D(A_1) \cap D(A_0^*) \Leftrightarrow H_1$ is compact, holds, if and only if $D(A_0) \Leftrightarrow H_0$, $D(A_1) \Leftrightarrow H_1$ are compact and $N_{0,1}$ is finite dimensional. Moreover, as Banach space adjoints we have*

$$H_0 \cong H_0' \Leftrightarrow D(A_0)' \Leftrightarrow D(A_0) \Leftrightarrow H_0 \Leftrightarrow D(A_0^*) \Leftrightarrow H_1 \Leftrightarrow H_1 \cong H_1' \Leftrightarrow D(A_0^*)',$$

and

$$H_1 \cong H_1' \Leftrightarrow D(A_1)' \Leftrightarrow D(A_1) \Leftrightarrow H_1 \Leftrightarrow D(A_1^*) \Leftrightarrow H_2 \Leftrightarrow H_2 \cong H_2' \Leftrightarrow D(A_1^*)'.$$

Especially, the assumption that $D(A_1) \cap D(A_0^) \Leftrightarrow H_1$ is compact, is equivalent to $\dim N_{0,1} < \infty$ and*

$$H_0 \Leftrightarrow D(A_0)', \quad H_2 \Leftrightarrow D(A_1^*)'$$

are compact. Thus, we observe that Theorem 4.7 is equivalent to [26, Theorem 2.5] and that the assumptions of Theorem 4.7 are practically equivalent to those of [26, Theorem 2.4].

5. APPLICATIONS

Whenever closed Hilbert complexes like (4.12) together with the corresponding compact embedding $D(A_1) \cap D(A_0^*) \Leftrightarrow H_1$ occur, we can apply the general A_0^* - A_1 -lemma, i.e., Theorem 4.7. In three dimensions we typically have three closed and densely defined linear operators A_0 , A_1 , and A_2 , satisfying the complex properties $R(A_0) \subset N(A_1)$ and $R(A_1) \subset N(A_2)$, i.e.,

$$(5.1) \quad \begin{array}{ccccccc} D(A_0) \subset H_0 & \xrightarrow{A_0} & D(A_1) \subset H_1 & \xrightarrow{A_1} & D(A_2) \subset H_2 & \xrightarrow{A_2} & H_3, \\ H_0 & \xleftarrow{A_0^*} & D(A_0^*) \subset H_1 & \xleftarrow{A_1^*} & D(A_1^*) \subset H_2 & \xleftarrow{A_2^*} & D(A_2^*) \subset H_3, \end{array}$$

together with the crucial compact embeddings

$$(5.2) \quad D(A_1) \cap D(A_0^*) \Leftrightarrow H_1, \quad D(A_2) \cap D(A_1^*) \Leftrightarrow H_2.$$

Let us recall our general assumptions on the underlying domain from Section 2: In the following examples we suppose that $\Omega \subset \mathbb{R}^3$ is a bounded weak Lipschitz domain, see [3, Definition 2.3], with boundary Γ , which is divided into two relatively open weak Lipschitz subsets Γ_t and $\Gamma_n := \Gamma \setminus \overline{\Gamma}_t$, see [3, Definition 2.5].

5.1. The div-rot-Lemma Revisited. The first and most canonical example is given by the classical operators from vector analysis

$$\begin{aligned} A_0 &:= \mathring{\nabla}_{\Gamma_t} : \mathring{H}_{\Gamma_t}^1(\Omega) \subset L^2(\Omega) \longrightarrow L_\epsilon^2(\Omega); & u &\mapsto \nabla u, \\ A_1 &:= \mu^{-1} \mathring{\text{rot}}_{\Gamma_t} : \mathring{R}_{\Gamma_t}(\Omega) \subset L_\epsilon^2(\Omega) \longrightarrow L_\mu^2(\Omega); & E &\mapsto \mu^{-1} \text{rot } E, \\ A_2 &:= \mathring{\text{div}}_{\Gamma_t} \mu : \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega) \subset L_\mu^2(\Omega) \longrightarrow L^2(\Omega); & H &\mapsto \text{div } \mu H. \end{aligned}$$

A_0 , A_1 , and A_2 are unbounded, densely defined, and closed linear operators with adjoints

$$\begin{aligned} A_0^* &= \mathring{\nabla}_{\Gamma_n}^* = -\mathring{\text{div}}_{\Gamma_n} \epsilon : \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \subset L_\epsilon^2(\Omega) \longrightarrow L^2(\Omega); & H &\mapsto -\text{div } \epsilon H, \\ A_1^* &= (\mu^{-1} \mathring{\text{rot}}_{\Gamma_t})^* = \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n} : \mathring{R}_{\Gamma_n}(\Omega) \subset L_\mu^2(\Omega) \longrightarrow L_\epsilon^2(\Omega); & E &\mapsto \epsilon^{-1} \text{rot } E, \\ A_2^* &= (\mathring{\text{div}}_{\Gamma_t} \mu)^* = -\mathring{\nabla}_{\Gamma_n} : \mathring{H}_{\Gamma_n}^1(\Omega) \subset L^2(\Omega) \longrightarrow L_\mu^2(\Omega); & u &\mapsto -\nabla u. \end{aligned}$$

Here, $\epsilon, \mu : \Omega \rightarrow \mathbb{R}^{3 \times 3}$ are symmetric and uniformly positive definite $L^\infty(\Omega)$ -tensor fields. The complex properties hold as

$$\begin{aligned} R(A_0) &= \mathring{\nabla}_{\Gamma_t} \mathring{H}_{\Gamma_t}^1(\Omega) \subset \mathring{R}_{\Gamma_t,0}(\Omega) = N(A_1), & R(A_1^*) &= \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n} \mathring{R}_{\Gamma_n}(\Omega) \subset \epsilon^{-1} \mathring{D}_{\Gamma_n,0}(\Omega) = N(A_0^*), \\ R(A_1) &= \mu^{-1} \mathring{\text{rot}}_{\Gamma_t} \mathring{R}_{\Gamma_t}(\Omega) \subset \mu^{-1} \mathring{D}_{\Gamma_t,0}(\Omega) = N(A_2), & R(A_2^*) &= \mathring{\nabla}_{\Gamma_n} \mathring{H}_{\Gamma_n}^1(\Omega) \subset \mathring{R}_{\Gamma_n,0}(\Omega) = N(A_1^*). \end{aligned}$$

Hence, the sequences (5.1) read

$$\begin{aligned} \mathring{H}_{\Gamma_t}^1(\Omega) \subset L^2(\Omega) &\xrightarrow{A_0 = \mathring{\nabla}_{\Gamma_t}} \mathring{R}_{\Gamma_t}(\Omega) \subset L_\epsilon^2(\Omega) \xrightarrow{A_1 = \mu^{-1} \mathring{\text{rot}}_{\Gamma_t}} \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega) \subset L_\mu^2(\Omega) \xrightarrow{A_2 = \mathring{\text{div}}_{\Gamma_t} \mu} L^2(\Omega), \\ L^2(\Omega) &\xleftarrow{A_0^* = -\mathring{\text{div}}_{\Gamma_n} \epsilon} \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \subset L_\epsilon^2(\Omega) \xleftarrow{A_1^* = \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n}} \mathring{R}_{\Gamma_n}(\Omega) \subset L_\mu^2(\Omega) \xleftarrow{A_2^* = -\mathring{\nabla}_{\Gamma_n}} \mathring{H}_{\Gamma_n}^1(\Omega) \subset L^2(\Omega). \end{aligned}$$

These are the well known Hilbert complexes for electro-magnetics, which are also known as de Rham complexes. Moreover, the crucial embeddings (5.2), i.e.,

$$D(A_1) \cap D(A_0^*) = \mathring{R}_{\Gamma_t}(\Omega) \cap \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega), \quad D(A_2) \cap D(A_1^*) = \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega) \cap \mathring{R}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega),$$

are compact by Weck's selection theorem (2.1), see [3, Theorem 4.7]. Indeed, Weck's selection theorems are independent of the material law tensors ϵ or μ . Choosing the pair (A_0, A_1) we get by Theorem 4.7 the following:

Theorem 5.1 (global div ϵ - μ^{-1} rot-lemma). *Let $(E_n) \subset \mathring{R}_{\Gamma_t}(\Omega)$ and $(H_n) \subset \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega)$ be two sequences bounded in $\mathring{R}(\Omega)$ resp. $\epsilon^{-1} \mathring{D}(\Omega)$. Then there exist $E \in \mathring{R}_{\Gamma_t}(\Omega)$ and $H \in \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega)$ as well as subsequences, again denoted by (E_n) and (H_n) , such that (E_n) and (H_n) converge weakly in $\mathring{R}(\Omega)$ resp. $\epsilon^{-1} \mathring{D}(\Omega)$ to E resp. H together with the convergence of the inner products*

$$\langle E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle E, H \rangle_{L^2(\Omega)}.$$

Remark 5.2. *We note:*

- (i) *Considering (E_n) and (ϵH_n) shows that Theorem 5.1 is equivalent to the global div-curl-lemma Theorem 3.1.*
- (ii) *Theorem 5.1 has a corresponding local version similar to the local div-curl-lemma Corollary 3.2 and Remark 3.3, which holds with no regularity or boundedness assumptions on Ω .*
- (iii) *Choosing the pair (A_1, A_2) we get by Theorem 4.7 a permutation of Theorem 5.1, shortly stating, that for bounded sequences $(E_n) \subset \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega)$ and $(H_n) \subset \mathring{R}_{\Gamma_n}(\Omega)$ it holds (after picking subsequences) $\langle E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle E, H \rangle_{L^2(\Omega)}$.*

Other examples are the following:

5.2. Generalized Electro-Magnetics. Let us allow $\Omega \subset \mathbb{R}^N$ or even to be a smooth Riemannian manifold with Lipschitz boundary and interface submanifolds Γ , and Γ_t, Γ_n . Using the calculus of alternating differential q -forms, we define the exterior derivative d and co-derivative $\delta = \pm * d *$ as Sobolev derivatives in the distributional weak sense by

$$D^q(\Omega) := \{E \in L^{2,q}(\Omega) : dE \in L^{2,q+1}(\Omega)\}, \quad \Delta^{q+1}(\Omega) := \{H \in L^{2,q+1}(\Omega) : \delta H \in L^{2,q}(\Omega)\}.$$

To introduce boundary conditions we define

$$\mathring{d}_{\Gamma_t}^q : \mathring{D}_{\Gamma_t}^q(\Omega) := \overline{\mathring{C}_{\Gamma_t}^{\infty,q}(\Omega)}^{D^q(\Omega)} \subset L^{2,q}(\Omega) \longrightarrow L^{2,q+1}(\Omega); \quad E \mapsto dE$$

as closure of the classical exterior derivative d acting on test q -forms. $\mathring{d}_{\Gamma_t}^q$ is an unbounded, densely defined, and closed linear operator with adjoint

$$(\mathring{d}_{\Gamma_t}^q)^* = -\mathring{\delta}_{\Gamma_n}^{q+1} : \mathring{D}_{\Gamma_n}^{q+1}(\Omega) := \overline{\mathring{C}_{\Gamma_n}^{\infty,q+1}(\Omega)}^{\Delta^{q+1}(\Omega)} \subset L^{2,q+1}(\Omega) \longrightarrow L^{2,q}(\Omega); \quad H \mapsto -\delta H.$$

Let us introduce

$$A_0 := \mathring{d}_{\Gamma_t}^{q-1}, \quad A_1 := \mathring{d}_{\Gamma_t}^q, \quad A_0^* = \mathring{\delta}_{\Gamma_n}^q, \quad A_1^* = \mathring{\delta}_{\Gamma_n}^{q+1}.$$

The complex properties hold as, e.g.,

$$R(A_0) = \mathring{d}_{\Gamma_t}^{q-1} \mathring{D}_{\Gamma_t}^{q-1}(\Omega) \subset \mathring{D}_{\Gamma_t,0}^q(\Omega) = N(A_1), \quad R(A_1^*) = \mathring{\delta}_{\Gamma_n}^{q+1} \mathring{\Delta}_{\Gamma_n}^{q+1}(\Omega) \subset \mathring{\Delta}_{\Gamma_n,0}^q(\Omega) = N(A_0^*)$$

by the classical properties $\delta \delta = \pm * d d * = 0$. Hence, the sequences (5.1) read

$$\begin{aligned} \mathring{D}_{\Gamma_t}^{q-1}(\Omega) \subset \mathbb{L}^{2,q-1}(\Omega) &\xrightarrow{A_0 = \mathring{d}_{\Gamma_t}^{q-1}} \mathring{D}_{\Gamma_t}^q(\Omega) \subset \mathbb{L}^{2,q}(\Omega) \xrightarrow{A_1 = \mathring{d}_{\Gamma_t}^q} \mathbb{L}^{2,q+1}(\Omega), \\ \mathbb{L}^{2,q-1}(\Omega) &\xleftarrow{A_0^* = \mathring{\delta}_{\Gamma_n}^q} \mathring{\Delta}_{\Gamma_n}^q(\Omega) \subset \mathbb{L}^{2,q}(\Omega) \xleftarrow{A_1^* = \mathring{\delta}_{\Gamma_n}^{q+1}} \mathring{\Delta}_{\Gamma_n}^{q+1}(\Omega) \subset \mathbb{L}^{2,q+1}(\Omega), \end{aligned}$$

which are the well known Hilbert complexes for generalized electro-magnetics, i.e., the de Rham complexes. Moreover, the crucial embedding (5.2), i.e.,

$$D(A_1) \cap D(A_0^*) = \mathring{D}_{\Gamma_t}^q(\Omega) \cap \mathring{\Delta}_{\Gamma_n}^q(\Omega) \hookrightarrow \mathbb{L}^2(\Omega),$$

is compact by a generalization of Weck's selection theorem (2.1), see [4] or the fundamental papers of Weck [28] and Picard [18] for full boundary conditions. Indeed, Weck's selection theorems are independent of the material law tensors ϵ or μ . Theorem 4.7 shows the following result:

Theorem 5.3 (global δ -d-lemma). *Let $(E_n) \subset \mathring{D}_{\Gamma_t}^q(\Omega)$ and $(H_n) \subset \mathring{\Delta}_{\Gamma_n}^q(\Omega)$ be two sequences bounded in $\mathbb{D}^q(\Omega)$ resp. $\mathbb{A}^q(\Omega)$. Then there exist $E \in \mathring{D}_{\Gamma_t}^q(\Omega)$ and $H \in \mathring{\Delta}_{\Gamma_n}^q(\Omega)$ as well as subsequences, again denoted by (E_n) and (H_n) , such that (E_n) and (H_n) converge weakly in $\mathbb{D}^q(\Omega)$ resp. $\mathbb{A}^q(\Omega)$ to E resp. H together with the convergence of the inner products*

$$\langle E_n, H_n \rangle_{\mathbb{L}^{2,q}(\Omega)} \rightarrow \langle E, H \rangle_{\mathbb{L}^{2,q}(\Omega)}.$$

Remark 5.4. *We note:*

- (i) For $N = 3$ and $q = 1$ (resp. $q = 2$) we obtain by Theorem 5.3 again the global div-curl-lemma Theorem 3.1.
- (ii) For $q = 0$ (resp. $q = N$) as well as identifying $\mathring{d}_{\Gamma_t}^0 = \mathring{\nabla}_{\Gamma_t}$ and $\mathring{\Delta}_{\Gamma_n}^0(\Omega) = 0$ (resp. $\mathring{d}_{\Gamma_t}^N = 0$ and $\mathring{\Delta}_{\Gamma_n}^N(\Omega) = \mathring{\nabla}_{\Gamma_n}$) we get by Theorem 5.3 the following trivial (by Rellich's selection theorem) result: For all bounded sequences $(u_n) \subset \mathring{H}_{\Gamma_t}^1(\Omega)$ and $(v_n) \subset \mathbb{L}^2(\Omega)$ there exist $u \in \mathring{H}_{\Gamma_t}^1(\Omega)$ and $v \in \mathbb{L}^2(\Omega)$ as well as subsequences, again denoted by (u_n) and (v_n) , such that (u_n) and (v_n) converge weakly in $\mathring{H}_{\Gamma_t}^1(\Omega)$ resp. $\mathbb{L}^2(\Omega)$ to u resp. v together with the convergence of the inner products $\langle u_n, v_n \rangle_{\mathbb{L}^2(\Omega)} \rightarrow \langle u, v \rangle_{\mathbb{L}^2(\Omega)}$.
- (iii) Theorem 5.3 has a corresponding local version similar to the local div-curl-lemma Corollary 3.2 and Remark 3.3, which holds with no regularity or boundedness assumptions on Ω .

5.3. Biharmonic Equation and General Relativity, Gravitational Waves. Let $\Omega \subset \mathbb{R}^3$ enjoy our general assumptions. We introduce symmetric and deviatoric (trace-free) square integrable tensor fields in $\mathbb{L}^2(\Omega; \mathbb{S})$ resp. $\mathbb{L}^2(\Omega; \mathbb{T})$ and as closures of the Hessian $\nabla \nabla$, and Rot, Div (row-wise rot, div), applied to test functions resp. test tensor fields, the linear operators

$$\begin{aligned} A_0 &:= \nabla \nabla_{\Gamma_t} : \mathring{H}_{\Gamma_t}^2(\Omega) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega)}^{\mathbb{H}^2(\Omega)} \subset \mathbb{L}^2(\Omega) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{S}); & u &\mapsto \nabla \nabla u, \\ A_1 &:= \text{Rot}_{\mathbb{S}, \Gamma_t} : \mathring{R}_{\Gamma_t}(\Omega; \mathbb{S}) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega; \mathbb{S})}^{\mathbb{R}(\Omega)} \subset \mathbb{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{T}); & S &\mapsto \text{Rot } S, \\ A_2 &:= \text{Div}_{\mathbb{T}, \Gamma_t} : \mathring{D}_{\Gamma_t}(\Omega; \mathbb{T}) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega; \mathbb{T})}^{\mathbb{D}(\Omega)} \subset \mathbb{L}^2(\Omega; \mathbb{T}) \longrightarrow \mathbb{L}^2(\Omega); & T &\mapsto \text{Div } T. \end{aligned}$$

A_0 , A_1 , and A_2 are unbounded, densely defined, and closed linear operators with adjoints

$$\begin{aligned} A_0^* &= (\nabla \nabla_{\Gamma_t})^* = \text{div Div}_{\mathbb{S}, \Gamma_n} : \mathring{D}D_{\Gamma_n}(\Omega; \mathbb{S}) := \overline{\mathring{C}_{\Gamma_n}^\infty(\Omega; \mathbb{S})}^{\mathbb{D}D(\Omega)} \subset \mathbb{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbb{L}^2(\Omega); & S &\mapsto \text{div Div } S, \\ A_1^* &= \text{Rot}_{\mathbb{S}, \Gamma_t}^* = \text{sym Rot}_{\mathbb{T}, \Gamma_n} : \mathring{R}_{\text{sym}, \Gamma_n}(\Omega; \mathbb{T}) := \overline{\mathring{C}_{\Gamma_n}^\infty(\Omega; \mathbb{T})}^{\mathbb{R}_{\text{sym}}(\Omega)} \subset \mathbb{L}^2(\Omega; \mathbb{T}) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{S}); & T &\mapsto \text{sym Rot } T, \\ A_2^* &= \text{Div}_{\mathbb{T}, \Gamma_t}^* = -\text{dev } \mathring{\nabla}_{\Gamma_n} : \mathring{H}_{\Gamma_n}^1(\Omega) \subset \mathbb{L}^2(\Omega) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{T}); & v &\mapsto -\text{dev } \nabla v, \end{aligned}$$

see [17] for details. Note that u , v , and S , T are scalar, vector, and tensor (matrix) fields, respectively. The complex properties hold as

$$\begin{aligned} R(A_0) &= \nabla^\circ \nabla_{\Gamma_t} \mathring{H}_{\Gamma_t}^2(\Omega) \subset \mathring{R}_{\Gamma_t,0}(\Omega; \mathbb{S}) = N(A_1), \\ R(A_1^*) &= \text{sym} \mathring{\text{Rot}}_{\mathbb{S}, \Gamma_n} \mathring{R}_{\text{sym}, \Gamma_n}(\Omega; \mathbb{T}) \subset \mathring{\text{DD}}_{\Gamma_n,0}(\Omega; \mathbb{S}) = N(A_0^*), \\ R(A_1) &= \mathring{\text{Rot}}_{\mathbb{S}, \Gamma_t} \mathring{R}_{\Gamma_t}(\Omega; \mathbb{S}) \subset \mathring{\text{D}}_{\Gamma_t,0}(\Omega; \mathbb{T}) = N(A_2), \\ R(A_2^*) &= \text{dev} \mathring{\nabla}_{\Gamma_n} \mathring{H}_{\Gamma_n}^1(\Omega) \subset \mathring{R}_{\text{sym}, \Gamma_n, 0}(\Omega; \mathbb{T}) = N(A_1^*). \end{aligned}$$

The sequences (5.1) read

$$\begin{aligned} \mathring{H}_{\Gamma_t}^2(\Omega) \subset L^2(\Omega) &\xrightarrow{A_0 = \nabla^\circ \nabla_{\Gamma_t}} \mathring{R}_{\Gamma_t}(\Omega; \mathbb{S}) \subset L^2(\Omega; \mathbb{S}) \xrightarrow{A_1 = \mathring{\text{Rot}}_{\mathbb{S}, \Gamma_t}} \mathring{\text{D}}_{\Gamma_t}(\Omega; \mathbb{T}) \subset L^2(\Omega; \mathbb{T}) \xrightarrow{A_2 = \text{Dev}_{\mathbb{T}, \Gamma_t}} L^2(\Omega), \\ L^2(\Omega) &\xleftarrow{A_0^* = \text{div} \mathring{\text{Div}}_{\mathbb{S}, \Gamma_n}} \mathring{\text{DD}}_{\Gamma_n}(\Omega; \mathbb{S}) \subset L^2(\Omega; \mathbb{S}) \xleftarrow{A_1^* = \text{sym} \mathring{\text{Rot}}_{\mathbb{T}, \Gamma_n}} \mathring{R}_{\text{sym}, \Gamma_n}(\Omega; \mathbb{T}) \subset L^2(\Omega; \mathbb{T}) \xleftarrow{A_2^* = -\text{dev} \mathring{\nabla}_{\Gamma_n}} \mathring{H}_{\Gamma_n}^1(\Omega) \subset L^2(\Omega). \end{aligned}$$

These are the so called Grad grad and div Div complexes, appearing, e.g., in the biharmonic problem or general relativity, see [17] for details. The crucial embeddings (5.2), i.e.,

$$\begin{aligned} D(A_1) \cap D(A_0^*) &= \mathring{R}_{\Gamma_t}(\Omega; \mathbb{S}) \cap \mathring{\text{DD}}_{\Gamma_n}(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S}), \\ D(A_2) \cap D(A_1^*) &= \mathring{\text{D}}_{\Gamma_t}(\Omega; \mathbb{T}) \cap \mathring{R}_{\text{sym}, \Gamma_n}(\Omega; \mathbb{T}) \hookrightarrow L^2(\Omega; \mathbb{T}), \end{aligned}$$

are compact by [17, Lemma 3.22]. Choosing the pair (A_0, A_1) we get by Theorem 4.7 the following:

Theorem 5.5 (global div Div-Rot- \mathbb{S} -lemma). *Let $(S_n) \subset \mathring{R}_{\Gamma_t}(\Omega; \mathbb{S})$ and $(T_n) \subset \mathring{\text{DD}}_{\Gamma_n}(\Omega; \mathbb{S})$ be two sequences bounded in $R(\Omega)$ resp. $\text{DD}(\Omega)$. Then there exist $S \in \mathring{R}_{\Gamma_t}(\Omega; \mathbb{S})$ and $T \in \mathring{\text{DD}}_{\Gamma_n}(\Omega; \mathbb{S})$ as well as subsequences, again denoted by (S_n) and (T_n) , such that (S_n) and (T_n) converge weakly in $R(\Omega)$ resp. $\text{DD}(\Omega)$ to S resp. T together with the convergence of the inner products*

$$\langle S_n, T_n \rangle_{L^2(\Omega, \mathbb{S})} \rightarrow \langle S, T \rangle_{L^2(\Omega, \mathbb{S})}.$$

For the pair (A_1, A_2) Theorem 4.7 implies:

Theorem 5.6 (global sym Rot-Div- \mathbb{T} -lemma). *Let $(S_n) \subset \mathring{\text{D}}_{\Gamma_t}(\Omega; \mathbb{T})$ and $(T_n) \subset \mathring{R}_{\text{sym}, \Gamma_n}(\Omega; \mathbb{T})$ be two sequences bounded in $\text{D}(\Omega)$ resp. $\mathring{R}_{\text{sym}}(\Omega)$. Then there exist $S \in \mathring{\text{D}}_{\Gamma_t}(\Omega; \mathbb{T})$ and $T \in \mathring{R}_{\text{sym}, \Gamma_n}(\Omega; \mathbb{T})$ as well as subsequences, again denoted by (S_n) and (T_n) , such that (S_n) and (T_n) converge weakly in $\text{D}(\Omega)$ resp. $\mathring{R}_{\text{sym}}(\Omega)$ to S resp. T together with the convergence of the inner products*

$$\langle S_n, T_n \rangle_{L^2(\Omega, \mathbb{T})} \rightarrow \langle S, T \rangle_{L^2(\Omega, \mathbb{T})}.$$

Remark 5.7. *Theorem 5.1 has a corresponding local version similar to the local div-curl-lemma Corollary 3.2 and Remark 3.3, which holds with no regularity or boundedness assumptions on Ω .*

5.4. Linear Elasticity. Let

$$\begin{aligned} A_0 &:= \text{sym} \mathring{\nabla}_{\Gamma_t} : \mathring{H}_{\Gamma_t}^1(\Omega) \subset L^2(\Omega) \longrightarrow L^2(\Omega; \mathbb{S}); & v &\mapsto \text{sym} \nabla v, \\ A_1 &:= \text{Rot} \mathring{\text{Rot}}_{\mathbb{S}, \Gamma_t}^\top : \mathring{\text{RR}}_{\Gamma_t}^\top(\Omega; \mathbb{S}) := \overline{\mathring{\text{C}}_{\Gamma_t}^\infty(\Omega; \mathbb{S})}^{\text{RR}^\top(\Omega)} \subset L^2(\Omega; \mathbb{S}) \longrightarrow L^2(\Omega; \mathbb{S}); & S &\mapsto \text{Rot} \text{Rot}^\top S, \\ A_2 &:= \mathring{\text{Div}}_{\mathbb{S}, \Gamma_t} : \mathring{\text{D}}_{\Gamma_t}(\Omega; \mathbb{S}) := \overline{\mathring{\text{C}}_{\Gamma_t}^\infty(\Omega; \mathbb{S})}^{\text{D}(\Omega)} \subset L^2(\Omega; \mathbb{S}) \longrightarrow L^2(\Omega); & T &\mapsto \text{Div} T. \end{aligned}$$

A_0 , A_1 , and A_2 are unbounded, densely defined, and closed linear operators with adjoints

$$\begin{aligned} A_0^* &= (\text{sym} \mathring{\nabla}_{\Gamma_t})^* = -\mathring{\text{Div}}_{\mathbb{S}, \Gamma_n} : \mathring{\text{D}}_{\Gamma_n}(\Omega; \mathbb{S}) \subset L^2(\Omega; \mathbb{S}) \longrightarrow L^2(\Omega); & S &\mapsto -\text{Div} S, \\ A_1^* &= (\text{Rot} \mathring{\text{Rot}}_{\mathbb{S}, \Gamma_t}^\top)^* = \text{Rot} \mathring{\text{Rot}}_{\mathbb{S}, \Gamma_n}^\top : \mathring{\text{RR}}_{\Gamma_n}^\top(\Omega; \mathbb{S}) \subset L^2(\Omega; \mathbb{S}) \longrightarrow L^2(\Omega; \mathbb{S}); & T &\mapsto \text{Rot} \text{Rot}^\top T, \\ A_2^* &= \mathring{\text{Div}}_{\mathbb{S}, \Gamma_t}^* = -\text{sym} \mathring{\nabla}_{\Gamma_n} : \mathring{H}_{\Gamma_n}^1(\Omega) \subset L^2(\Omega) \longrightarrow L^2(\Omega; \mathbb{S}); & v &\mapsto -\text{sym} \nabla v. \end{aligned}$$

Note that v resp. S, T are vector resp. tensor (matrix) fields. The complex properties hold as

$$\begin{aligned} R(A_0) &= \text{sym } \overset{\circ}{\nabla}_{\Gamma_t} \overset{\circ}{\mathbb{H}}_{\Gamma_t}^1(\Omega) \subset \overset{\circ}{\text{RR}}_{\Gamma_t,0}^\top(\Omega; \mathbb{S}) = N(A_1), \\ R(A_1^*) &= \text{Rot } \overset{\circ}{\text{Rot}}_{\mathbb{S}, \Gamma_n}^\top \overset{\circ}{\text{RR}}_{\Gamma_n}^\top(\Omega; \mathbb{S}) \subset \overset{\circ}{\mathbb{D}}_{\Gamma_n,0}(\Omega; \mathbb{S}) = N(A_0^*), \\ R(A_1) &= \text{Rot } \overset{\circ}{\text{Rot}}_{\mathbb{S}, \Gamma_t}^\top \overset{\circ}{\text{RR}}_{\Gamma_t}^\top(\Omega; \mathbb{S}) \subset \overset{\circ}{\mathbb{D}}_{\Gamma_t,0}(\Omega; \mathbb{S}) = N(A_2), \\ R(A_2^*) &= \text{sym } \overset{\circ}{\nabla}_{\Gamma_n} \overset{\circ}{\mathbb{H}}_{\Gamma_n}^1(\Omega) \subset \overset{\circ}{\text{RR}}_{\Gamma_n,0}^\top(\Omega; \mathbb{S}) = N(A_1^*). \end{aligned}$$

The sequences (5.1) read

$$\begin{aligned} \overset{\circ}{\mathbb{H}}_{\Gamma_t}^1(\Omega) \subset \text{L}^2(\Omega) &\xrightarrow{A_0 = \text{sym } \overset{\circ}{\nabla}_{\Gamma_t}} \overset{\circ}{\text{RR}}_{\Gamma_t}^\top(\Omega; \mathbb{S}) \subset \text{L}^2(\Omega; \mathbb{S}) \xrightarrow{A_1 = \text{Rot } \overset{\circ}{\text{Rot}}_{\mathbb{S}, \Gamma_t}^\top} \overset{\circ}{\mathbb{D}}_{\Gamma_t}(\Omega; \mathbb{S}) \subset \text{L}^2(\Omega; \mathbb{S}) \xrightarrow{A_2 = \text{Div}_{\mathbb{S}, \Gamma_t}} \text{L}^2(\Omega), \\ \text{L}^2(\Omega) &\xleftarrow{A_0^* = -\text{Div}_{\mathbb{S}, \Gamma_n}} \overset{\circ}{\mathbb{D}}_{\Gamma_n}(\Omega; \mathbb{S}) \subset \text{L}^2(\Omega; \mathbb{S}) \xleftarrow{A_1^* = \text{Rot } \overset{\circ}{\text{Rot}}_{\mathbb{S}, \Gamma_n}^\top} \overset{\circ}{\text{RR}}_{\Gamma_n}^\top(\Omega; \mathbb{S}) \subset \text{L}^2(\Omega; \mathbb{S}) \xleftarrow{A_2^* = -\text{sym } \overset{\circ}{\nabla}_{\Gamma_n}} \overset{\circ}{\mathbb{H}}_{\Gamma_n}^1(\Omega) \subset \text{L}^2(\Omega). \end{aligned}$$

These are the so called Rot Rot complexes, appearing, e.g., in linear elasticity, see [17]. The crucial embeddings (5.2), i.e.,

$$\begin{aligned} D(A_1) \cap D(A_0^*) &= \overset{\circ}{\text{RR}}_{\Gamma_t}^\top(\Omega; \mathbb{S}) \cap \overset{\circ}{\mathbb{D}}_{\Gamma_n}(\Omega; \mathbb{S}) \hookrightarrow \text{L}^2(\Omega; \mathbb{S}), \\ D(A_2) \cap D(A_1^*) &= \overset{\circ}{\mathbb{D}}_{\Gamma_t}(\Omega; \mathbb{S}) \cap \overset{\circ}{\text{RR}}_{\Gamma_n}^\top(\Omega; \mathbb{S}) \hookrightarrow \text{L}^2(\Omega; \mathbb{S}), \end{aligned}$$

are compact, which can be proved by the same techniques showing [17, Lemma 3.22]. Choosing the pair (A_0, A_1) we get by Theorem 4.7 the following:

Theorem 5.8 (global Rot Rot[⊤]-Div- \mathbb{S} -lemma). *Let $(S_n) \subset \overset{\circ}{\text{RR}}_{\Gamma_t}^\top(\Omega; \mathbb{S})$ and $(T_n) \subset \overset{\circ}{\mathbb{D}}_{\Gamma_n}(\Omega; \mathbb{S})$ be two sequences bounded in $\overset{\circ}{\text{RR}}^\top(\Omega)$ resp. $\overset{\circ}{\mathbb{D}}(\Omega)$. Then there exist $S \in \overset{\circ}{\text{RR}}_{\Gamma_t}^\top(\Omega; \mathbb{S})$ and $T \in \overset{\circ}{\mathbb{D}}_{\Gamma_n}(\Omega; \mathbb{S})$ as well as subsequences, again denoted by (S_n) and (T_n) , such that (S_n) and (T_n) converge weakly in $\overset{\circ}{\text{RR}}^\top(\Omega)$ resp. $\overset{\circ}{\mathbb{D}}(\Omega)$ to S resp. T together with the convergence of the inner products*

$$\langle S_n, T_n \rangle_{\text{L}^2(\Omega, \mathbb{S})} \rightarrow \langle S, T \rangle_{\text{L}^2(\Omega, \mathbb{S})}.$$

Remark 5.9. *Let us note:*

- (i) *Theorem 4.7 for the pair (A_1, A_2) implies the same result just interchanging S_n, T_n and the boundary conditions.*
- (ii) *Theorem 5.8 has a corresponding local version similar to the local div-curl-lemma Corollary 3.2 and Remark 3.3, which holds with no regularity or boundedness assumptions on Ω .*
- (iii) *We emphasize the strong symmetry in the Rot Rot complexes of linear elasticity.*

REFERENCES

- [1] A. Alexanderian. Expository paper: a primer on homogenization of elliptic pdes with stationary and ergodic random coefficient functions. *Rocky Mountain J. Math.*, 45(3):703–735, 2015.
- [2] S. Bartels. Numerical analysis of a finite element scheme for the approximation of harmonic maps into surfaces. *Math. Comp.*, 79(271):1263–1301, 2010.
- [3] S. Bauer, D. Pauly, and M. Schomburg. The Maxwell compactness property in bounded weak Lipschitz domains with mixed boundary conditions. *SIAM J. Math. Anal.*, 48(4):2912–2943, 2016.
- [4] S. Bauer, D. Pauly, and M. Schomburg. Weck’s selection theorem for bounded weak Lipschitz domains and mixed boundary conditions. *submitted*, 2017.
- [5] M. Briane, J. Casado-Dáz, and F. Murat. The div-curl lemma “trente ans après”: an extension and an application to the g-convergence of unbounded monotone operators. *J. Math. Pures Appl.*, (9) 91(5), 2009.
- [6] M. Costabel. A remark on the regularity of solutions of Maxwell’s equations on Lipschitz domains. *Math. Methods Appl. Sci.*, 12(4):365–368, 1990.
- [7] L.C. Evans. *Weak convergence methods for nonlinear partial differential equations*. American Mathematical Society, Providence, 1990.
- [8] L.C. Evans. Partial regularity for stationary harmonic maps into spheres. *Arch. Rational Mech. Anal.*, 116(2):101–113, 1991.
- [9] A. Freire, S. Müller, and M. Struwe. Weak compactness of wave maps and harmonic maps. *Ann. Inst. H. Poincaré Anal.*, Non Linéaire 15(6):725–754, 1998.

- [10] A. Gloria, S. Neukamm, and F. Otto. Quantification of ergodicity in stochastic homogenization: optimal bounds via spectral gap on glauher dynamics. *Invent. Math.*, 199(2):455–515, 2015.
- [11] F. Jochmann. A compactness result for vector fields with divergence and curl in $L^q(\Omega)$ involving mixed boundary conditions. *Appl. Anal.*, 66:189–203, 1997.
- [12] R. Leis. *Initial Boundary Value Problems in Mathematical Physics*. Teubner, Stuttgart, 1986.
- [13] F. Murat. Compacité par compensation. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 5(3):489–507, 1978.
- [14] D. Pauly. On constants in Maxwell inequalities for bounded and convex domains. *Zapiski POMI*, 435:46–54, 2014, & *J. Math. Sci. (N.Y.)*, 210(6):787–792, 2015.
- [15] D. Pauly. On Maxwell’s and Poincaré’s constants. *Discrete Contin. Dyn. Syst. Ser. S*, 8(3):607–618, 2015.
- [16] D. Pauly. On the Maxwell constants in 3D. *Math. Methods Appl. Sci.*, 40(2):435–447, 2017.
- [17] D. Pauly and W. Zulehner. On closed and exact Grad-grad- and div-Div-complexes, corresponding compact embeddings for tensor rotations, and a related decomposition result for biharmonic problems in 3D. <https://arxiv.org/abs/1609.05873>, submitted, 2017.
- [18] R. Picard. An elementary proof for a compact imbedding result in generalized electromagnetic theory. *Math. Z.*, 187:151–164, 1984.
- [19] T. Rivière. Conservation laws for conformally invariant variational problems. *Invent. Math.*, 168(1):1–22, 2007.
- [20] B. Schweizer. On friedrichs inequality, helmholtz decomposition, vector potentials, and the div-curl lemma. *preprint*, <http://www.mathematik.uni-dortmund.de/lisi/schweizer/Preprints/publist.html>, 2017.
- [21] B. Schweizer and M. Röger. Strain gradient visco-plasticity with dislocation densities contributing to the energy. *preprint*, <http://www.mathematik.uni-dortmund.de/lisi/schweizer/Preprints/publist.html>, <https://arxiv.org/abs/1704.05326>, 2017.
- [22] M. Struwe. *Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems*. Springer, Berlin, 2008.
- [23] L. Tartar. Compensated compactness and applications to partial differential equations. *Nonlinear analysis and mechanics, Heriot-Watt symposium*, 4:136–211, 1979.
- [24] L. Tartar. *The general theory of homogenization. A personalized introduction*. Springer, Berlin, 2009.
- [25] L. Tartar. Compensated compactness with more geometry. *Springer Proc. Math. Stat.*, 137:74–101, 2015.
- [26] M. Waurick. A functional analytic perspective to the div-curl lemma. *preprint*, <https://arxiv.org/abs/1703.09593>, 2017.
- [27] C. Weber. A local compactness theorem for Maxwell’s equations. *Math. Methods Appl. Sci.*, 2:12–25, 1980.
- [28] N. Weck. Maxwell’s boundary value problems on Riemannian manifolds with nonsmooth boundaries. *J. Math. Anal. Appl.*, 46:410–437, 1974.
- [29] K.-J. Witsch. A remark on a compactness result in electromagnetic theory. *Math. Methods Appl. Sci.*, 16:123–129, 1993.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, CAMPUS ESSEN, GERMANY
E-mail address, Dirk Pauly: dirk.pauly@uni-due.de

IN DER SCHRIFTENREIHE DER FAKULTÄT FÜR MATHEMATIK ZULETZT ERSCHIENENE
BEITRÄGE:

- Nr. 769: Mali, O., Muzalevskiy, A., Pauly, D.: Conforming and Non-Conforming Functional A Posteriori Error Estimates for Elliptic Boundary Value Problems in Exterior Domains: Theory and Numerical Tests, 2013
- Nr. 770: Bauer, S., Neff, P., Pauly, D., Starke, G.: Dev-Div- and DevSym-DevCurl-Inequalities for Incompatible Square Tensor Fields with Mixed Boundary Conditions, 2013
- Nr. 771: Pauly, D.: On the Maxwell Inequalities for Bounded and Convex Domains, 2013
- Nr. 772: Pauly, D.: On Maxwell's and Poincaré's Constants, 2013
- Nr. 773: Fried, M. N., Jahnke, H. N.: Otto Toeplitz's "The problem of university infinitesimal calculus courses and their demarcation from infinitesimal calculus in high schools" (1927), 2013
- Nr. 774: Yurko, V.: Spectral Analysis for Differential Operators of Variable Orders on Star-type Graphs: General Case, 2014
- Nr. 775: Freiling, G., Yurko, V.: Differential Operators on Hedgehog-type Graphs with General Matching Conditions, 2014
- Nr. 776: Anjam, I., Pauly, D.: Functional A Posteriori Error Equalities for Conforming Mixed Approximations of Elliptic Problems, 2014
- Nr. 777: Pauly, D.: On the Maxwell Constants in 3D, 2014
- Nr. 778: Pozzi, P.: Computational Anisotropic Willmore Flow, 2014
- Nr. 779: Buterin, S.A., Freiling, G., Yurko, V.A.: Lectures on the Theory of entire Functions, 2014
- Nr. 780: Blatt, S., Reiter, Ph.: Modeling repulsive forces on fibres via knot energies, 2014
- Nr. 781: Neff, P., Ghiba, I.-D., Lankeit, J.: The exponentiated Hencky-logarithmic strain energy. Part I: Constitutive issues and rank-one convexity, 2014
- Nr. 782: Neff, P., Münch, I., Martin, R.: Rediscovering G.F. Becker's early axiomatic deduction of a multiaxial nonlinear stress-strain relation based on logarithmic strain, 2014
- Nr. 783: Neff, P., Ghiba, I.-D., Madeo, A., Placidi, L., Rosi, G.: A unifying perspective: the relaxed linear micromorphic continuum, 2014
- Nr. 784: Müller, F.: On $C^{1,1/2}$ -regularity of H -surfaces with a free boundary, 2014
- Nr. 785: Müller, F.: Projectability of stable, partially free H -surfaces in the non-perpendicular case, 2015
- Nr. 786: Bauer S., Pauly, D.: On Korn's First Inequality for Tangential or Normal Boundary Conditions with Explicit Constants, 2015
- Nr. 787: Neff, P., Eidel, B., Martin, R.J.: Geometry of logarithmic strain measures in solid mechanics, 2015
- Nr. 788: Borisov, L., Neff, P., Sra, S., Thiel, Chr.: The sum of squared logarithms inequality in arbitrary dimensions, 2015
- Nr. 789: Bauer, S., Pauly, D., Schomburg, M.: The Maxwell Compactness Property in Bounded Weak Lipschitz Domains with Mixed Boundary Conditions, 2015
- Nr. 790: Claus, M., Krätschmer, V., Schultz, R.: WEAK CONTINUITY OF RISK FUNCTIONALS WITH APPLICATIONS TO STOCHASTIC PROGRAMMING, 2015
- Nr. 790a: Claus, M., Krätschmer, V., Schultz, R.: WEAK CONTINUITY OF RISK FUNCTIONALS WITH APPLICATIONS TO STOCHASTIC PROGRAMMING (Revision), 2015/2016
- Nr. 791: Bauer, S., Pauly, D.: On Korn's First Inequality for Mixed Tangential and Normal Boundary Conditions on Bounded Lipschitz-Domains in \mathbb{R}^N , 2016
- Nr. 792: Anjam, I., Pauly, D.: Functional A Posteriori Error Control for Conforming Mixed Approximations of Coercive Problems with Lower Order Terms, 2016
- Nr. 793: Herkenrath, U.: "ARS CONJECTANDI" UND DIE NATUR DES ZUFALLS, 2016

- Nr. 794: Martin, R. J., Ghiba, I.-D., Neff, P.: Rank-one convexity implies polyconvexity for isotropic, objective and isochoric elastic energies in the two-dimensional case, 2016
- Nr. 795: Fischle, A., Neff, P.: The geometrically nonlinear Cosserat micropolar shear-stretch energy. Part I: A general parameter reduction formula and energy-minimizing microrotations in 2D, 2016
- Nr. 796: Münch, I., Neff, P., Madeo, A., Ghiba, I.-D.: The modified indeterminate couple stress model: Why Yang et al.'s arguments motivating a symmetric couple stress tensor contain a gap and why the couple stress tensor may be chosen symmetric nevertheless, 2016
- Nr. 797: Madeo, A., Ghiba, I.-D., Neff, P., Münch, I.: A new view on boundary conditions in the Grioli-Koiter-Mindlin-Toupin indeterminate couple stress model, 2016
- Nr. 798: Claus, M.: ON STABILITY IN RISK AVERSE STOCHASTIC BILEVEL PROGRAMMING, 2016
- Nr. 799: Burtscheidt, J., Claus, M.: A Note on Stability for Risk Averse Stochastic Complementarity Problems, 2016
- Nr. 800: Pauly, D., Picard, R.: A Note on the Justification of the Eddy Current Model in Electrodynamics, 2016
- Nr. 801: Pauly, D., Yousept, I.: A Posteriori Error Analysis for the Optimal Control of Magneto-Static Fields, 2016
- Nr. 802: Zimmermann, A.: Martingale solutions for a pseudomonotone evolution equation with multiplicative noise, 2016
- Nr. 803: Tennstädt, T.: MEAN CONVEXITY OF THE ZERO SET OF SYMMETRIC MINIMAL SURFACES, 2016
- Nr. 804: Tennstädt, T.: HÖLDER CONTINUITY FOR CONTINUOUS SOLUTIONS OF THE SINGULAR MINIMAL SURFACE EQUATION WITH ARBITRARY ZERO SET, 2016
- Nr. 805: Pauly, D., Zulehner, W.: On Closed and Exact Grad grad- and div Div-Complexes, Corresponding Compact Embeddings for Symmetric Rotations, and a Related Decomposition Result for Biharmonic Problems in 3D, 2016
- Nr. 806: Dierkes, U., Tennstädt, T.: BERNSTEIN RESULTS FOR SYMMETRIC MINIMAL SURFACES OF CONTROLLED GROWTH, 2016
- Nr. 807: Wittbold, P., Scholtes, M.: Existence of Entropy Solutions to a Doubly Nonlinear Integro-Differential Equation, 2016
- Nr. 808: Zimmermann, A.: On a pseudomonotone evolution equation with multiplicative noise, 2016
- Nr. 809: Pauly, D.: Solution Theory and Functional A Posteriori Error Estimates for General First Order Systems with Applications to Electro-Magneto-Statics, 2016
- Nr. 810: Anjam, I., Pauly, D.: An Elementary Method of Deriving A Posteriori Error Equalities and Estimates for Linear Partial Differential Equations, 2016
- Nr. 811: Pauly, D.: On the Maxwell and Friedrichs/Poincaré Constants in ND, 2017
- Nr. 812: Pauly, D.: A Global div-curl-Lemma for Mixed Boundary Conditions in Weak Lipschitz Domains and a Corresponding Generalized A^*_0 - A_1 -Lemma in Hilbert Spaces, 2017