

Gelfand's trick for the spherical derived Hecke algebra

Lennart Gehrman

September 27th, 2020

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Motivation

Γ “nice” arithmetic group, e.g.:

- $\Gamma = \mathrm{SL}_2(\mathbb{Z}), \mathrm{SL}_3(\mathbb{Z}), \dots$
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$, K imaginary quadratic
- congruence subgroups, e.g., $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$

Γ acts on a symmetric space, e.g.:

- $\mathrm{SL}_2(\mathbb{Z})$ acts on the complex upper half plane
- $\Gamma = \mathrm{SL}_2(\mathcal{O}_K)$ acts on hyperbolic 3-space

Hecke algebra acts on $H^i(\Gamma \backslash X, \mathbb{Q})$

Phenomenon: same Hecke eigensystem can occur in several degrees

Venkatesh \rightsquigarrow global derived Hecke algebra $\mathcal{H}_{\mathbb{Q}_\ell}^*$:

- $\mathcal{H}_{\mathbb{Q}_\ell}^*$ is a graded \mathbb{Q}_ℓ -algebra
- $H^*(\Gamma \backslash X, \mathbb{Q}_\ell)$ is a graded $\mathcal{H}_{\mathbb{Q}_\ell}^*$ -module

Conjecture: derived Hecke algebra “causes” the phenomenon

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Setup

F/\mathbb{Q}_p finite extension, \mathcal{O} ring of integers, $\mathcal{O}/(\varpi) = \mathbb{F}_q$

\mathbb{G}/\mathcal{O} split reductive group

$\rightsquigarrow G = \mathbb{G}(F)$, $K = \mathbb{G}(\mathcal{O})$, $T \subseteq G$ maximal split torus

Example: $\mathbb{G} = \mathrm{GL}_n$

$\rightsquigarrow G = \mathrm{GL}_n(F)$, $K = \mathrm{GL}_n(\mathcal{O})$, $T =$ invertible diagonal matrices

Cartan decomposition (Bruhat-Tits): $G = KTK$

If $G = \mathrm{GL}_n(F)$, this is just the elementary divisor theorem

Classical spherical Hecke algebra

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Classical spherical Hecke algebra

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Classical spherical Hecke algebra

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Classical spherical Hecke algebra

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Classical spherical Hecke algebra

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Spherical Hecke algebra of G :

$$\mathcal{H}(G)_{\mathbb{C}} = \mathbb{C}[K \backslash G / K] = C_c(K \backslash G / K, \mathbb{C})$$

Product given by convolution:

$$(h_1 * h_2)(g) = \int_G h_1(gx^{-1}) \cdot h_2(x) dx$$

$G = KTK \Rightarrow \{\mathbb{1}_{KtK} \mid t \in T\}$ generates $\mathcal{H}(G)_{\mathbb{C}}$

Gelfand's trick $\Rightarrow \mathcal{H}(G)_{\mathbb{C}}$ is commutative

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Gelfand's trick for the spherical Hecke algebra

$\exists \sigma: G \rightarrow G$ involution such that

- $\sigma(K) = K$
- $\sigma(t) = t^{-1} \quad \forall t \in T$

If $G = \mathrm{GL}_n(F)$, take $\sigma(g) = {}^t g^{-1}$

Define $\sigma: \mathcal{H}(G)_{\mathbb{C}} \rightarrow \mathcal{H}(G)_{\mathbb{C}}$ by

$$\sigma(f)(g) = f(\sigma(g)^{-1})$$

Easy calculation $\rightsquigarrow \sigma(h_1 * h_2) = \sigma(h_2) * \sigma(h_1)$

Cartan decomposition $\Rightarrow \sigma(h) = h$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G/K]) \\ &= \text{Hom}_G(\mathbb{C}[G/K], \mathbb{C}[G/K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G/K], \mathbb{Z}/\ell^r\mathbb{Z}[G/K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G/K]) \\ &= \text{Hom}_G(\mathbb{C}[G/K], \mathbb{C}[G/K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G/K], \mathbb{Z}/\ell^r\mathbb{Z}[G/K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G/K]) \\ &= \text{Hom}_G(\mathbb{C}[G/K], \mathbb{C}[G/K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G/K], \mathbb{Z}/\ell^r\mathbb{Z}[G/K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G/K]) \\ &= \text{Hom}_G(\mathbb{C}[G/K], \mathbb{C}[G/K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G/K], \mathbb{Z}/\ell^r\mathbb{Z}[G/K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G/K]) \\ &= \text{Hom}_G(\mathbb{C}[G/K], \mathbb{C}[G/K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G/K], \mathbb{C}[G/K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G/K], \mathbb{Z}/\ell^r\mathbb{Z}[G/K])$$

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G / K]) \\ &= \text{Hom}_G(\mathbb{C}[G / K], \mathbb{C}[G / K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G / K], \mathbb{Z}/\ell^r\mathbb{Z}[G / K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G / K]) \\ &= \text{Hom}_G(\mathbb{C}[G / K], \mathbb{C}[G / K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G / K], \mathbb{Z}/\ell^r\mathbb{Z}[G / K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G / K]) \\ &= \text{Hom}_G(\mathbb{C}[G / K], \mathbb{C}[G / K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G / K], \mathbb{Z}/\ell^r\mathbb{Z}[G / K])$$

Derived Hecke algebra

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G / K]) \\ &= \text{Hom}_G(\mathbb{C}[G / K], \mathbb{C}[G / K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G / K], \mathbb{Z}/\ell^r\mathbb{Z}[G / K])$$

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G / K]) \\ &= \text{Hom}_G(\mathbb{C}[G / K], \mathbb{C}[G / K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G / K], \mathbb{Z}/\ell^r\mathbb{Z}[G / K])$$

$$\begin{aligned}\mathcal{H}(G)_{\mathbb{C}} &= \mathbb{C}[K \backslash G / K] \\ &= \text{Hom}_K(\mathbb{C}, \mathbb{C}[G / K]) \\ &= \text{Hom}_G(\mathbb{C}[G / K], \mathbb{C}[G / K])\end{aligned}$$

Idea:

$$\mathcal{H}^*(G)_{\mathbb{C}} := \bigoplus_i \text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K])$$

Problem:

$$\text{Ext}_G^i(\mathbb{C}[G / K], \mathbb{C}[G / K]) = 0 \text{ for } i \geq 1$$

Solution: choose $\ell \neq p$ and consider

$$\mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}} := \bigoplus_i \text{Ext}_G^i(\mathbb{Z}/\ell^r\mathbb{Z}[G / K], \mathbb{Z}/\ell^r\mathbb{Z}[G / K])$$

The main result

The main result

Theorem

$\ell \neq p$ prime

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$ and
- ℓ does not divide the order of the Weyl group of G

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$ and
- ℓ does not divide the order of the Weyl group of G

$\Rightarrow \mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}}$ is graded-commutative

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$ and
- ℓ does not divide the order of the Weyl group of G

$\Rightarrow \mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}}$ is graded-commutative for all r

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$ and
- ℓ does not divide the order of the Weyl group of G

$\Rightarrow \mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}}$ is graded-commutative for all r

Weyl group of GL_n isomorphic to S_n

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$ and
- ℓ does not divide the order of the Weyl group of G

$\Rightarrow \mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}}$ is graded-commutative for all r

Weyl group of GL_n isomorphic to S_n

\rightsquigarrow second condition equivalent to $\ell > n$ if $G = \mathrm{GL}_n(F)$

The main result

Theorem

$\ell \neq p$ prime such that

- $q \equiv 1 \pmod{\ell}$ and
- ℓ does not divide the order of the Weyl group of G

$\Rightarrow \mathcal{H}^*(G)_{\mathbb{Z}/\ell^r\mathbb{Z}}$ is graded-commutative for all r

Weyl group of GL_n isomorphic to S_n

\rightsquigarrow second condition equivalent to $\ell > n$ if $G = \mathrm{GL}_n(F)$

Result previously known under the condition $q \equiv 1 \pmod{\ell^r}$
(Venkatesh)

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

An explicit model for the derived Hecke algebra

$(x, y) \in G/K \times G/K \rightsquigarrow G_{xy}$ its stabilizer in G

An element $h \in \mathcal{H}(G)_R^*$ is a collection of elements

$$h(x, y) \in H^*(G_{xy}, R), \quad (x, y) \in G/K \times G/K$$

such that

- h is G -invariant, i.e. $\text{Ad}(g)^* h(gx, gy) = h(x, y)$
- h has finite support modulo G

Multiplication given by convolution:

$$(h_1 * h_2)(x, z) = \sum_{y \in G_{xz} \backslash G/K} \text{Cores}_{G_{xyz}}^{G_{xz}} (\text{Res}_{G_{xyz}}^{G_{xy}} h_1(x, y) \cup \text{Res}_{G_{xyz}}^{G_{yz}} h_2(y, z))$$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:
 $\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:
 $\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

The involution

$G=KTK \Rightarrow (x, y)$ and $(\sigma(y), \sigma(x))$ lie in same G -orbit, i.e.:

$\exists g_{xy} \in G$ such that $g_{xy}(x, y) = (\sigma(y), \sigma(x))$

$$g_{xy} G_{xy} g_{xy}^{-1} = \sigma(G_{xy})$$

$$\rightsquigarrow \text{Ad}(g_{xy})^* \sigma^* : H^*(G_{xy}, R) \rightarrow H^*(G_{xy}, R)$$

Define $h^\sigma \in \mathcal{H}(G)_R^*$ via $h^\sigma(x, y) = \text{Ad}(g_{xy})^* \sigma^*(h(x, y))$

Easy calculation

$$\rightsquigarrow \sigma(h_1 * h_2) = (-1)^{ij} \cdot \sigma(h_2) * \sigma(h_1), \deg(h_1) = i, \deg(h_2) = j$$

Under assumptions of theorem: $\sigma(h_1 * h_2) = \sigma(h_1) * \sigma(h_2)$

How to prove it?

How to prove it?

Would you like to know more?

How to prove it?

Would you like to know more?
Check out my preprint!