

# On exceptional zeros for $GL(2)$

Lennart Gehrman

June 10th, 2020

# Structure of talk

- 1 Introduction
- 2 P-adic L-functions
- 3 Exceptional zeroes

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM$ ,  $p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in \mathcal{S}_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in \mathcal{S}_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$



# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Mazur–Tate–Teitelbaum conjecture

$f = \sum_n a_n q^n \in S_{2k}(N)$  newform,  $N = pM, p \nmid M$   
 $\rightsquigarrow L_p(f, s)$   $p$ -adic  $L$ -function

$$L_p(f, k) \approx (1 - p^{k-1}/a_p) \cdot L(f, k)$$

$a_p = p^{k-1} \rightsquigarrow$  exceptional zero

Conjecture (Mazur–Tate–Teitelbaum):

$$\frac{d}{ds} L_p(f, s)|_{s=k} \approx \mathcal{L}(f) \cdot L(f, k)$$

$\mathcal{L}(f)$  is a local invariant, i.e.  $\mathcal{L}(f)$  only depends on the restriction of  $p$ -adic Galois representation attached to  $f$  to a decomposition group at  $p$

# Tate periods

$2k = 2$ ,  $a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM$ ,  $p \nmid M$ ,  $a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

# Tate periods

$2k = 2$ ,  $a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM$ ,  $p \nmid M$ ,  $a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

# Tate periods

$2k = 2, a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM, p \nmid M, a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

# Tate periods

$2k = 2, a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM, p \nmid M, a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

# Tate periods

$2k = 2, a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM, p \nmid M, a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$



# Tate periods

$2k = 2, a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM, p \nmid M, a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

# Tate periods

$2k = 2$ ,  $a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM$ ,  $p \nmid M$ ,  $a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

## Theorem (Tate)

# Tate periods

$2k = 2, a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM, p \nmid M, a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

## Theorem (Tate)

$\text{a} \quad q \in \mathbb{Q}_p, 0 < |q|_p < 1$

# Tate periods

$2k = 2, a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM, p \nmid M, a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

## Theorem (Tate)

$q \in \mathbb{Q}_p, 0 < |q|_p < 1$

$\rightsquigarrow E_q/\mathbb{Q}_p$  such that  $E_q(\mathbb{C}_p) \cong \mathbb{C}_p^*/q^{\mathbb{Z}}$

# Tate periods

$2k = 2$ ,  $a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM$ ,  $p \nmid M$ ,  $a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

## Theorem (Tate)

a  $q \in \mathbb{Q}_p$ ,  $0 < |q|_p < 1$

$\rightsquigarrow E_q/\mathbb{Q}_p$  such that  $E_q(\mathbb{C}_p) \cong \mathbb{C}_p^*/q^{\mathbb{Z}}$

b If  $E/\mathbb{Q}_p$  has split multiplicative reduction

# Tate periods

$2k = 2$ ,  $a_n \in \mathbb{Q}$  for all  $n$

$f \longleftrightarrow$  elliptic curve  $E_f/\mathbb{Q}$

$N = pM$ ,  $p \nmid M$ ,  $a_p = 1$

$\Rightarrow E_f$  has split multiplicative reduction at  $p$

## Theorem (Tate)

a  $q \in \mathbb{Q}_p$ ,  $0 < |q|_p < 1$

$\rightsquigarrow E_q/\mathbb{Q}_p$  such that  $E_q(\mathbb{C}_p) \cong \mathbb{C}_p^*/q^{\mathbb{Z}}$

b If  $E/\mathbb{Q}_p$  has split multiplicative reduction

$\rightsquigarrow E \cong E_q$  for a unique  $q \in \mathbb{Q}_p$ ,  $0 < |q|_p < 1$

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$

$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens

What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$

$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens

What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita



# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$

$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens

What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$
$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens  
What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$
$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens

What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$
$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens  
What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$
$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens  
What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$
$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens  
What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# $\mathcal{L}$ -invariants

$$\mathcal{L}(E_q) = \frac{\log_p(q)}{\text{ord}_p(q)} \rightsquigarrow \text{isogeny invariant}$$
$$\mathcal{L}(f) = \mathcal{L}(E_{f, \mathbb{Q}_p})$$

MTT conjecture proven by Greenberg–Stevens  
What about higher weights?

- Definition of  $\mathcal{L}$ -invariants by Fontaine–Mazur, Teitelbaum, Coleman, Darmon, Orton, Breuil
- Proof of MTT by Kato–Kurihara–Tsuji, Stevens, Darmon, Orton, Emerton
- Equality of  $\mathcal{L}$ -invariants by Coleman–Iovita, Iovita–Spieß, Breuil, Bertolini–Darmon–Iovita

# GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_\rho(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_\rho(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
 multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams



## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams



## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomena:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

## GL<sub>2</sub> over other number fields

$F$  number field,  $\pi$  cohomological automorphic representation of  $GL_2/F$

small slope condition  $\rightsquigarrow L_p(f, \underline{s})$  Barrera-Williams

new phenomenons:

- function in many variables if  $F$  is not totally real  
 $\rightsquigarrow$  need to compute partial derivatives in several directions
- $L_p(f, k) \approx \prod_{p|p} e_p(k) \cdot L(f, k)$   
multiple  $e_p(k) = 0 \rightsquigarrow$  compute higher partial derivatives

Known cases:

- $F$  totally real, parallel weight 2, first order: Mok
- $F$  totally real, parallel weight 2: Spieß
- $F$  arbitrary, parallel weight 2: Deppe, Bergunde
- $F$  totally real: Barrera–Dimitrov–Jorza
- $F$  imaginary quadratic, first order: Barrera–Williams

- 1 Introduction
- 2 P-adic L-functions
- 3 Exceptional zeroes



# $\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \twoheadrightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p = 2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \longrightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \longmapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

# $\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \twoheadrightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p = 2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \longrightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \longmapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

# $\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p = 2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \mapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

$\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p=2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \longrightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \longmapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

$\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p=2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \mapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

$\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \twoheadrightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p=2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \longrightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \longmapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

$\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \twoheadrightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p=2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \longrightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \longmapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$

# $\mathbb{Z}_p$ -extensions

$F$  number field,  $F_\infty/F$  extension with  $\mathcal{G} = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$

Example:  $\text{Gal}(F(\mu_{p^\infty})/F) \cong \mathbb{Z}_p \times \text{finite}$

$F_\infty$  is unramified outside  $p$

put  $F_p = F \otimes \mathbb{Q}_p \cong \prod_{p|p} F_p$ ,  $\mathcal{O}_{F,S_p} = \mathcal{O}_F[1/p]$

$\rightsquigarrow$  class field theory:

$$\text{rec}: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow \mathcal{G}$$

Fix isomorphism  $\ell: \text{Gal}(F_\infty/F) \xrightarrow{\cong} p\mathbb{Z}_p$  (resp.  $4\mathbb{Z}_2$  if  $p = 2$ )

$$\chi_\ell: F_p^*/\mathcal{O}_{F,S_p}^* \longrightarrow C^{\text{an}}(\mathbb{Z}_p, \mathbb{Z}_p)^*$$

$$\gamma \longmapsto [s \mapsto \exp_p(s \cdot \ell(\text{rec}(\gamma)))]$$



# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathfrak{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathfrak{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathfrak{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathfrak{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathfrak{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathfrak{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, s) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, s) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$



# Homology classes

$R$  a  $\mathbb{Q}_p$ -algebra,  $\chi: F_p^*/\mathcal{O}_{F,S_p}^* \rightarrow R^*$  locally analytic character

$\rightsquigarrow \chi \in H^0(\mathcal{O}_{F,S_p}^*, C^{\text{an}}(F_p^*, R))$

$E \subseteq \mathcal{O}_F^*$  torsion-free, finite index

Dirichlet's unit theorem  $\Rightarrow E \cong \mathbb{Z}^r$ ,  $r = r_{\mathbb{R}} + r_{\mathbb{C}} - 1$

$$\Rightarrow H_r(E, \mathbb{Z}) \cong \mathbb{Z}$$

Shapiro  $\rightsquigarrow$  fundamental class  $\vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c(F_p^*/U_p, \mathbb{Z}))$

$$c_{\chi} = \chi \cap \vartheta \in H_r(\mathcal{O}_{F,S_p}^*, C_c^{\text{an}}(F_p^*, R))$$

Given  $\kappa \in H^r(\mathcal{O}_{F,S_p}^*, \text{Dist}(F_p^*, E))$ ,  $E/\mathbb{Q}_p$  finite

$$\rightsquigarrow L_p(\kappa, \ell, \mathbf{s}) = \kappa \cap \chi_{\ell} \in C^{\text{an}}(\mathbb{Z}_p, E)$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^V)_{\mathfrak{m}_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{r_{\mathbb{R}}} \times \mathbb{H}_3^{r_{\mathbb{C}}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$\mathfrak{m}_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^V)_{\mathfrak{m}_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x x \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^V))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$\mathfrak{m}_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x X \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{r_{\mathbb{R}}} \times \mathbb{H}_3^{r_{\mathbb{C}}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$m_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x X \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{m_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$m_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x X \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{m_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$m_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x X \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{m_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$m_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{m_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x X \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{m_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$\mathfrak{m}_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x X \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$



# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$\mathfrak{m}_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x x \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$\mathfrak{m}_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x x \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Cohomology of locally symmetric spaces

$\pi$  cohomological cuspidal automorphic representation of  $\mathrm{PGL}_{2,F}$

$$\Rightarrow H_c^*(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi} \neq 0$$

$\Gamma_0(\mathfrak{n}) \subseteq \mathrm{PGL}_2(\mathcal{O}_F)$  congruence subgroup

$X = \mathbb{H}_2^{\mathbb{R}} \times \mathbb{H}_3^{\mathbb{C}}$  symmetric space

$V_k$  algebraic representation of  $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$

$\mathfrak{m}_\pi$  ideal of Hecke algebra given by  $\pi$

Moreover:

$$\dim H_c^{r+1}(\Gamma_0(\mathfrak{n}) \backslash X, V_{\underline{k}}^\vee)_{\mathfrak{m}_\pi}^\epsilon = 1$$

Put  $\Delta_0 = \ker(\mathbb{Z}[\mathbb{P}^1(F)] \rightarrow \mathbb{Z}, \sum_x n_x x \mapsto \sum_x n_x)$

$$\rightsquigarrow H^i(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, M)) \xrightarrow{\cong} H_c^{i+1}(\Gamma_0(\mathfrak{n}) \backslash X, M)$$

$$\Rightarrow H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Overconvergent cohomology

$I_p = \prod_{\mathfrak{p}|\rho} I_{\mathfrak{p}} \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_{\mathfrak{p}} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid \mathfrak{p} \text{ divides } c \right\}$$

$$\mathcal{A}_{\underline{k}} = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^{\underline{k}} f(g) \right\}$$

Have embedding  $V_{\underline{k}} \hookrightarrow \mathcal{A}_{\underline{k}}$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^f(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

# Overconvergent cohomology

$I_p = \prod_{\mathfrak{p}|\rho} I_{\mathfrak{p}} \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_{\mathfrak{p}} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid \mathfrak{p} \text{ divides } c \right\}$$

$$\mathcal{A}_{\underline{k}} = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^k f(g) \right\}$$

Have embedding  $V_{\underline{k}} \hookrightarrow \mathcal{A}_{\underline{k}}$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^f(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

# Overconvergent cohomology

$I_p = \prod_{\mathfrak{p}|\rho} I_{\mathfrak{p}} \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_{\mathfrak{p}} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid \mathfrak{p} \text{ divides } c \right\}$$

$$\mathcal{A}_{\underline{k}} = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^k f(g) \right\}$$

Have embedding  $V_{\underline{k}} \hookrightarrow \mathcal{A}_{\underline{k}}$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^f(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

# Overconvergent cohomology

$I_p = \prod_{p|p} I_p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_p = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid p \text{ divides } c \right\}$$

$$\mathcal{A}_k = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^k f(g) \right\}$$

Have embedding  $V_k \hookrightarrow \mathcal{A}_k$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_k^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_k^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_k^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Overconvergent cohomology

$I_p = \prod_{\mathfrak{p}|\rho} I_{\mathfrak{p}} \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_{\mathfrak{p}} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid \mathfrak{p} \text{ divides } c \right\}$$

$$\mathcal{A}_{\underline{k}} = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^{\underline{k}} f(g) \right\}$$

Have embedding  $V_{\underline{k}} \hookrightarrow \mathcal{A}_{\underline{k}}$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^f(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$



# Overconvergent cohomology

$I_p = \prod_{p|p} I_p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_p = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid p \text{ divides } c \right\}$$

$$\mathcal{A}_k = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^k f(g) \right\}$$

Have embedding  $V_k \hookrightarrow \mathcal{A}_k$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_k^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_k^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_k^\vee))_{\mathfrak{m}_\pi}^\epsilon = 1$$

# Overconvergent cohomology

$I_p = \prod_{\mathfrak{p}|\rho} I_{\mathfrak{p}} \subseteq \mathrm{PGL}_2(\mathcal{O}_{F_p})$  Iwahori subgroup

$$I_{\mathfrak{p}} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PGL}_2(\mathcal{O}_{F_p}) \mid \mathfrak{p} \text{ divides } c \right\}$$

$$\mathcal{A}_{\underline{k}} = \left\{ f: I_p \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = (d/a)^{\underline{k}} f(g) \right\}$$

Have embedding  $V_{\underline{k}} \hookrightarrow \mathcal{A}_{\underline{k}}$

$$\rightsquigarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \rightarrow H^i(\Gamma_0(n), \mathrm{Hom}(\Delta_0, V_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

Ash-Stevens: This is an isomorphism for small slopes. In particular:

$$H^f(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

# Cohomology of $p$ -arithmetic groups

$\Gamma_0(n)^p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F,S_p})$   $p$ -arithmetic congruence subgroup

$$\rightsquigarrow H^r(\Gamma_0(n)^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^c \xrightarrow{\cong} H^r(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^c$$

$I(\chi)$  locally analytic principal series:

$$I(\chi) = \left\{ f: G(F_p) \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = \chi(d/a) f(g) \right\}$$

$\chi: F_p^* \rightarrow E^*$ ,  $\chi(x) = x^k$  if  $x$  integral,  $\chi(\varpi) = U_p$ -eigenvalue

Proof via Koszul resolution by Kohlhaase–Schraen

# Cohomology of $p$ -arithmetic groups

$\Gamma_0(\mathfrak{n})^p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F,S_p})$   $p$ -arithmetic congruence subgroup

$$\rightsquigarrow H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \xrightarrow{\cong} H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

$I(\chi)$  locally analytic principal series:

$$I(\chi) = \left\{ f: G(F_p) \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = \chi(d/a) f(g) \right\}$$

$\chi: F_p^* \rightarrow E^*$ ,  $\chi(x) = x^k$  if  $x$  integral,  $\chi(\varpi) = U_p$ -eigenvalue

Proof via Koszul resolution by Kohlhaase–Schraen

# Cohomology of $p$ -arithmetic groups

$\Gamma_0(\mathfrak{n})^p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F,S_p})$   $p$ -arithmetic congruence subgroup

$$\rightsquigarrow H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \xrightarrow{\cong} H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

$I(\chi)$  locally analytic principal series:

$$I(\chi) = \left\{ f: G(F_p) \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = \chi(d/a) f(g) \right\}$$

$\chi: F_p^* \rightarrow E^*$ ,  $\chi(x) = x^k$  if  $x$  integral,  $\chi(\varpi) = U_p$ -eigenvalue

Proof via Koszul resolution by Kohlhaase–Schraen

# Cohomology of $p$ -arithmetic groups

$\Gamma_0(\mathfrak{n})^p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F,S_p})$   $p$ -arithmetic congruence subgroup

$$\rightsquigarrow H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \xrightarrow{\cong} H^r(\Gamma_0(\mathfrak{n}), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

$I(\chi)$  locally analytic principal series:

$$I(\chi) = \left\{ f: G(F_p) \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = \chi(d/a) f(g) \right\}$$

$\chi: F_p^* \rightarrow E^*$ ,  $\chi(x) = x^k$  if  $x$  integral,  $\chi(\varpi) = U_p$ -eigenvalue  
 Proof via Koszul resolution by Kohlhaase–Schraen

# Cohomology of $p$ -arithmetic groups

$\Gamma_0(n)^p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F,S_p})$   $p$ -arithmetic congruence subgroup

$$\rightsquigarrow H^r(\Gamma_0(n)^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \xrightarrow{\cong} H^r(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

$I(\chi)$  locally analytic principal series:

$$I(\chi) = \left\{ f: G(F_p) \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = \chi(d/a) f(g) \right\}$$

$\chi: F_p^* \rightarrow E^*$ ,  $\chi(x) = x^k$  if  $x$  integral,  $\chi(\varpi) = U_p$ -eigenvalue

Proof via Koszul resolution by Kohlhaase–Schraen

# Cohomology of $p$ -arithmetic groups

$\Gamma_0(n)^p \subseteq \mathrm{PGL}_2(\mathcal{O}_{F,S_p})$   $p$ -arithmetic congruence subgroup

$$\rightsquigarrow H^r(\Gamma_0(n)^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \xrightarrow{\cong} H^r(\Gamma_0(n), \mathrm{Hom}(\Delta_0, \mathcal{A}_{\underline{k}}^\vee))_{\mathfrak{m}_\pi}^\epsilon$$

$I(\chi)$  locally analytic principal series:

$$I(\chi) = \left\{ f: G(F_p) \rightarrow E \text{ locally analytic} \mid f \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g \right) = \chi(d/a) f(g) \right\}$$

$\chi: F_p^* \rightarrow E^*$ ,  $\chi(x) = x^k$  if  $x$  integral,  $\chi(\varpi) = U_p$ -eigenvalue

Proof via Koszul resolution by Kohlhaase–Schraen



# The map $\delta$

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, S_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

# The map $\delta$

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, S_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

The map  $\delta$ 

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, \mathfrak{S}_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

The map  $\delta$ 

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, \mathfrak{S}_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

# The map $\delta$

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, \mathfrak{S}_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

# The map $\delta$

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, \mathfrak{S}_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

# The map $\delta$

Embed  $F_p^*$  in  $\mathrm{PGL}_2(F_p)$  via  $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

Want  $F_p^*$ -equivariant map

$$\delta: C_c^{\mathrm{an}}(F_p^*, E) \rightarrow I(\chi)$$

$$\delta^*: H^r(\Gamma_0(\mathfrak{n})^p, \mathrm{Hom}(\Delta_0, I(\chi)^\vee))_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^r(\mathcal{O}_{F, S_p}^*, \mathrm{Dist}(F_p^*, E))$$

$$\rightsquigarrow \kappa_\pi \rightsquigarrow L_p(\pi, \ell, \mathfrak{s}) = L_p(\kappa_\pi, \ell, \mathfrak{s})$$

$\delta$  is given by trivialization of  $I(\chi)$  on open subset of  $\mathbb{P}^1$ :

$$\delta(f)(g) = f(u)\chi(u)\chi(d/a) \text{ for } g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$$

- 1 Introduction
- 2 P-adic L-functions
- 3 Exceptional zeroes



# The map $\delta$ again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in S_{\text{sp}} \subseteq S_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in S_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin S_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, S_{\text{sp}}}^*, \text{Dist}(\prod_{\mathfrak{p} \in S_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin S_{\text{sp}}} F_{\mathfrak{p}}^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{s=0} L_{\rho}(\pi, \ell, s) \geq |S_{\text{sp}}|$   
 + cohomological formula for  $|S_{\text{sp}}|$ -th derivative

# The map $\delta$ again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \subseteq \mathcal{S}_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, \mathcal{S}_{\text{sp}}}^*, \text{Dist}(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{\mathfrak{s}=0} L_{\rho}(\pi, \ell, \mathfrak{s}) \geq |\mathcal{S}_{\text{sp}}|$   
 + cohomological formula for  $|\mathcal{S}_{\text{sp}}|$ -th derivative

# The map $\delta$ again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \subseteq \mathcal{S}_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, \mathcal{S}_{\text{sp}}}^*, \text{Dist}(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{\mathfrak{s}=0} L_{\rho}(\pi, \ell, \mathfrak{s}) \geq |\mathcal{S}_{\text{sp}}|$   
 + cohomological formula for  $|\mathcal{S}_{\text{sp}}|$ -th derivative

# The map $\delta$ again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \subseteq \mathcal{S}_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, \mathcal{S}_{\text{sp}}}^*, \text{Dist}(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E))$$

Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{s=0} L_{\rho}(\pi, \ell, s) \geq |\mathcal{S}_{\text{sp}}|$   
 + cohomological formula for  $|\mathcal{S}_{\text{sp}}|$ -th derivative

# The map $\delta$ again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \subseteq \mathcal{S}_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, \mathcal{S}_{\rho}}^*, \text{Dist}(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{\mathfrak{s}=0} L_{\rho}(\pi, \ell, \mathfrak{s}) \geq |\mathcal{S}_{\text{sp}}|$   
 + cohomological formula for  $|\mathcal{S}_{\text{sp}}|$ -th derivative

# The map $\delta$ again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \subseteq \mathcal{S}_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, \mathcal{S}_{\rho}}^*, \text{Dist}(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{s=0} L_{\rho}(\pi, \ell, \mathbf{s}) \geq |\mathcal{S}_{\text{sp}}|$   
 + cohomological formula for  $|\mathcal{S}_{\text{sp}}|$ -th derivative

## The map $\delta$ again

$\chi = \prod_{p|\rho} \chi_p$  with  $\chi_p: F_p^* \rightarrow E^*$

Suppose  $\chi_p$  can be extended to analytic function on  $F_p$  for  $p \in S_{sp} \subseteq S_\rho$

$$\delta: C_c^{\text{an}}\left(\prod_{p \in S_{sp}} F_p \times \prod_{p \notin S_{sp}} F_p^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_\pi \in H^r(\mathcal{O}_{F, S_{sp}}^*, \text{Dist}(\prod_{p \in S_{sp}} F_p \times \prod_{p \notin S_{sp}} F_p^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{s=0} L_\rho(\pi, \ell, s) \geq |S_{sp}|$   
 + cohomological formula for  $|S_{sp}|$ -th derivative

### Lemma

$\chi_p$  can be extended to analytic function on  $F_p$

The map  $\delta$  again

$\chi = \prod_{\mathfrak{p}|\rho} \chi_{\mathfrak{p}}$  with  $\chi_{\mathfrak{p}}: F_{\mathfrak{p}}^* \rightarrow E^*$

Suppose  $\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$  for  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \subseteq \mathcal{S}_{\rho}$

$$\delta: C_c^{\text{an}}\left(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E\right) \rightarrow I(\chi)$$

$\Rightarrow \kappa_{\pi} \in H^r(\mathcal{O}_{F, \mathcal{S}_{\text{sp}}}^*, \text{Dist}(\prod_{\mathfrak{p} \in \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}} \times \prod_{\mathfrak{p} \notin \mathcal{S}_{\text{sp}}} F_{\mathfrak{p}}^*, E))$   
 Spieß, Dasgupta-Spiess  $\rightsquigarrow \text{ord}_{s=0} L_{\rho}(\pi, \ell, s) \geq |\mathcal{S}_{\text{sp}}|$   
 + cohomological formula for  $|\mathcal{S}_{\text{sp}}|$ -th derivative

## Lemma

$\chi_{\mathfrak{p}}$  can be extended to analytic function on  $F_{\mathfrak{p}}$   
 if and only if  $\chi_{\mathfrak{p}}(x) = x^{k_{\mathfrak{p}}}$  for all  $x$



# The Theorem

## Theorem

Let  $n = |S_{\text{sp}}|$ .

$$\frac{d^n}{ds^n} L(\pi, \ell, s)|_{s=0} \approx \prod_{p \in S_{\text{sp}}} \mathcal{L}_{\ell}^{\text{aut}}(\pi, p) \cdot \prod_{p \notin S_{\text{sp}}} e_p(1/2) \cdot L(\pi, 1/2)$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\text{lore}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \rightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\text{lore}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \rightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\text{lore}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \rightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\ell \text{orec}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

$$(i): \mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \rightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

$$(ii): \text{Breuil} \rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\ell \text{orec}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \twoheadrightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\ell \text{orec}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \twoheadrightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\ell \text{orec}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \twoheadrightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(n)^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$



# Automorphic $\mathcal{L}$ -invariants I

Let  $\mathfrak{p} \in \mathcal{S}_{\text{sp}}$ . Want to construct:

$$\text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow E, \lambda \mapsto \mathcal{L}_{\lambda}(\pi, \mathfrak{p})$$

such that  $\mathcal{L}_{\text{ord}_{\mathfrak{p}}}(\pi, \mathfrak{p}) = 1$

$$\rightsquigarrow \mathcal{L}_{\ell}^{\text{aut}}(\pi, \mathfrak{p}) = \mathcal{L}_{\ell \text{orec}_{\mathfrak{p}}}^{\text{aut}}(\pi, \mathfrak{p})$$

Two steps:

(i):  $\mathfrak{p} \in \mathcal{S}_{\text{sp}} \Rightarrow I(\chi_{\mathfrak{p}}) \twoheadrightarrow \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}}$

$$\rightsquigarrow \dim H^r(\Gamma_0(\mathfrak{n})^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} = 1$$

(ii): Breuil  $\rightsquigarrow \text{Hom}(F_{\mathfrak{p}}^*, E) \rightarrow \text{Ext}_{\text{PGL}_2(F_{\mathfrak{p}})}^1(V_{\underline{k}_{\mathfrak{p}}}, \text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})$

$$H^r(\Gamma_0(\mathfrak{n})^{\mathfrak{p}}, \text{Hom}(\Delta_0, (\text{St}(\underline{k}_{\mathfrak{p}})^{\text{an}})^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon} \xrightarrow{\cup \lambda} H^{r+1}(\Gamma_0(\mathfrak{n})^{\mathfrak{p}}, \text{Hom}(\Delta_0, V_{\underline{k}_{\mathfrak{p}}}^{\vee}))_{\mathfrak{m}_{\pi}}^{\epsilon}$$

# Automorphic $\mathcal{L}$ -invariants II

## Lemma

*The cup product with the  $p$ -adic valuation  $\text{ord}_p$  is an isomorphism.*

# Automorphic $\mathcal{L}$ -invariants II

## Lemma

*The cup product with the  $p$ -adic valuation  $\text{ord}_p$  is an isomorphism.*

## Definition

$$\mathcal{L}_\lambda^{\text{aut}}(\pi, \mathfrak{p})^\epsilon = (\cup \text{ord}_p)^{-1} \circ (\cup \lambda)$$