

Big principal series and L-invariants

Lennart Gehrmann

June 30th, 2020

Structure of talk

- 1 Motivation: p -adic periods of elliptic curves
- 2 Darmon's L-invariant
- 3 Big principal series and p -adic families

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Elliptic curves

k field, $\text{char}(k) \neq 2, 3$

E/k elliptic curve, $E = V(Y^2 - X^3 - aX - b)$ with $a, b \in k$
such that $4a^3 + 27b^2 \neq 0$

Alternatively: E/k connected, smooth, projective curve of
genus 1 with k -rational point

Alternatively: E/k one-dimensional, connected, projective,
algebraic group

Complex uniformization

$$k = \mathbb{C}$$

$\rightsquigarrow E(\mathbb{C})$ is a complex one-dimensional torus, i.e.:

$$E(\mathbb{C}) \cong \mathbb{C}/\Lambda, \quad \Lambda \subseteq \mathbb{C} \text{ lattice}$$

$$\text{Wlog } \Lambda = \mathbb{Z} + \tau\mathbb{Z}, \quad \Im(\tau) > 0$$

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\exp(2\pi i \cdot)} \mathbb{C}^*$$

yields

$$E(\mathbb{C}) \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z} \cong \mathbb{C}^*/q^{\mathbb{Z}}, \quad 0 < |q| < 1$$

Complex uniformization

$$k = \mathbb{C}$$

$\rightsquigarrow E(\mathbb{C})$ is a complex one-dimensional torus, i.e.:

$$E(\mathbb{C}) \cong \mathbb{C}/\Lambda, \Lambda \subseteq \mathbb{C} \text{ lattice}$$

$$\text{Wlog } \Lambda = \mathbb{Z} + \tau\mathbb{Z}, \Im(\tau) > 0$$

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\exp(2\pi i \cdot)} \mathbb{C}^*$$

yields

$$E(\mathbb{C}) \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z} \cong \mathbb{C}^*/q^{\mathbb{Z}}, 0 < |q| < 1$$

Complex uniformization

$$k = \mathbb{C}$$

$\rightsquigarrow E(\mathbb{C})$ is a complex one-dimensional torus, i.e.:

$$E(\mathbb{C}) \cong \mathbb{C}/\Lambda, \Lambda \subseteq \mathbb{C} \text{ lattice}$$

$$\text{Wlog } \Lambda = \mathbb{Z} + \tau\mathbb{Z}, \Im(\tau) > 0$$

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\exp(2\pi i \cdot)} \mathbb{C}^*$$

yields

$$E(\mathbb{C}) \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z} \cong \mathbb{C}^*/q^{\mathbb{Z}}, 0 < |q| < 1$$

Complex uniformization

$$k = \mathbb{C}$$

$\rightsquigarrow E(\mathbb{C})$ is a complex one-dimensional torus, i.e.:

$$E(\mathbb{C}) \cong \mathbb{C}/\Lambda, \Lambda \subseteq \mathbb{C} \text{ lattice}$$

$$\text{Wlog } \Lambda = \mathbb{Z} + \tau\mathbb{Z}, \Im(\tau) > 0$$

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\exp(2\pi i \cdot)} \mathbb{C}^*$$

yields

$$E(\mathbb{C}) \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z} \cong \mathbb{C}^*/q^{\mathbb{Z}}, 0 < |q| < 1$$

Complex uniformization

$$k = \mathbb{C}$$

$\rightsquigarrow E(\mathbb{C})$ is a complex one-dimensional torus, i.e.:

$$E(\mathbb{C}) \cong \mathbb{C}/\Lambda, \Lambda \subseteq \mathbb{C} \text{ lattice}$$

$$\text{Wlog } \Lambda = \mathbb{Z} + \tau\mathbb{Z}, \Im(\tau) > 0$$

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\exp(2\pi i \cdot)} \mathbb{C}^*$$

yields

$$E(\mathbb{C}) \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z} \cong \mathbb{C}^*/q^{\mathbb{Z}}, 0 < |q| < 1$$

Complex uniformization

$$k = \mathbb{C}$$

$\rightsquigarrow E(\mathbb{C})$ is a complex one-dimensional torus, i.e.:

$$E(\mathbb{C}) \cong \mathbb{C}/\Lambda, \Lambda \subseteq \mathbb{C} \text{ lattice}$$

$$\text{Wlog } \Lambda = \mathbb{Z} + \tau\mathbb{Z}, \Im(\tau) > 0$$

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\exp(2\pi i \cdot)} \mathbb{C}^*$$

yields

$$E(\mathbb{C}) \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z} \cong \mathbb{C}^*/q^{\mathbb{Z}}, 0 < |q| < 1$$

P-adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

P-adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

P-adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

Theorem (Tate 1959)

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

Theorem (Tate 1959)

$q \in \mathbb{Q}_p$, $0 < |q|_p < 1$

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

Theorem (Tate 1959)

$q \in \mathbb{Q}_p$, $0 < |q|_p < 1$

$\rightsquigarrow E_q/\mathbb{Q}_p$ such that $E_q(\mathbb{Q}_p) \cong \mathbb{Q}_p^*/q^{\mathbb{Z}}$

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

Theorem (Tate 1959)

- a $q \in \mathbb{Q}_p$, $0 < |q|_p < 1$
 $\rightsquigarrow E_q/\mathbb{Q}_p$ such that $E_q(\mathbb{Q}_p) \cong \mathbb{Q}_p^*/q^{\mathbb{Z}}$
- b *If E/\mathbb{Q}_p has split multiplicative reduction*

p -adic uniformization

What about $k = \mathbb{Q}_p$?

$\Lambda \subseteq \mathbb{Q}_p$ discrete $\rightsquigarrow \Lambda = \{0\}$

But: $q \in \mathbb{Q}_p$, $0 < |q|_p < 1 \rightsquigarrow q^{\mathbb{Z}} \subseteq \mathbb{Q}_p^*$ discrete, cocompact

Theorem (Tate 1959)

- a $q \in \mathbb{Q}_p$, $0 < |q|_p < 1$
 $\rightsquigarrow E_q/\mathbb{Q}_p$ such that $E_q(\mathbb{Q}_p) \cong \mathbb{Q}_p^*/q^{\mathbb{Z}}$
- b If E/\mathbb{Q}_p has split multiplicative reduction
 $\rightsquigarrow E \cong E_q$, $q \in \mathbb{Q}_p$, $0 < |q|_p < 1$

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface
 \rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p
(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Modularity

$$k = \mathbb{Q}$$

(Wiles et al.) $\rightsquigarrow \exists f \in \mathcal{S}_2(\Gamma_0(N))$ such that

$$L(E, s) = L(f, s)$$

f is a differential form on a Riemann surface

\rightsquigarrow purely analytic object

Suppose E has split multiplicative reduction at prime p

(Tate) $\rightsquigarrow q_E \in \mathbb{Q}_p^*$

Question: Can we recover q_E from f ?

Arithmetic L-Invariants

Answer:

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

Arithmetic L-Invariants

Answer: No!

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

Arithmetic L-Invariants

Answer: No!

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

Arithmetic L-Invariants

Answer: No!

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

Arithmetic L-Invariants

Answer: No!

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

Arithmetic L-Invariants

Answer: No!

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

Arithmetic L-Invariants

Answer: No!

$E \sim E'$ isogenous $\rightsquigarrow L(E, s) = L(E', s)$

but: $q_E^n = q_{E'}^m$ for $m, n \in \mathbb{Z}_{>0}$

$$\rightsquigarrow \mathcal{L}_p(E) = \frac{\log_p(q_E)}{\text{ord}_p(q_E)} \text{ isogeny invariant}$$

Question: Can we recover $\mathcal{L}_p(E)$ from f ?

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

Theorem (Bertolini-Darmon-Iovita 2010)

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

Theorem (Bertolini-Darmon-Iovita 2010)

$$\mathcal{L}^D(f) = -2a'_p$$

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

Theorem (Bertolini-Darmon-Iovita 2010)

$$\mathcal{L}^D(f) = -2a'_p$$

Aim: give new proof

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

Theorem (Bertolini-Darmon-Iovita 2010)

$$\mathcal{L}^D(f) = -2a'_p$$

Aim: give new proof that generalizes to higher rank groups

A zoo of L-Invariants

Answer: Yes!

Teitelbaum: $\mathcal{L}^T(f)$ using Cerednik-Drinfeld uniformization

Darmon: $\mathcal{L}^D(f)$ using cohomology of $SL_2(\mathbb{Z}[1/p])$

Breuil: $\mathcal{L}^B(f)$ using completed cohomology

Greenberg-Stevens: $a'_p =$ derivative of U_p -eigenvalue of Hida family through f

Theorem (Bertolini-Darmon-Iovita 2010)

$$\mathcal{L}^D(f) = -2a'_p$$

Aim: give new proof that generalizes to higher rank groups
(joint work with Giovanni Rosso)

- 1 Motivation: p -adic periods of elliptic curves
- 2 Darmon's L-invariant
- 3 Big principal series and p -adic families

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 1: Cohomology of arithmetic groups

$E \longleftrightarrow f$ weight 2 modular form of level N

$$\Gamma_0(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$$

$\rightsquigarrow f$ (holomorphic) 1-form on $\Gamma_0(N) \backslash \mathbb{H}$

$$f \in H^1(\Gamma_0(N) \backslash \mathbb{H}, \mathbb{C}) \cong H^1(\Gamma_0(N), \mathbb{C})$$

Multiplicity one theorem:

$$\dim H^1(\Gamma_0(N), \mathbb{C})^+[f] = 1$$

$$\dim H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f] = 1$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{\text{constant}\}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+[f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{ \text{constant} \}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+[f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{\text{constant}\}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+[f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{\text{constant}\}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+ [f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+ [f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{ \text{constant} \}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+[f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{\text{constant}\}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+[f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 2: Cohomology of p-arithmetic groups

E split multiplicative reduction at p

$$\Gamma_0^p(N) = \left\{ \gamma \in \mathrm{GL}_2(\mathbb{Z}[1/p]) \mid \gamma \cong \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}, \det(\gamma) > 0 \right\}$$

$$\mathrm{St}_p(R) = \left\{ \phi: \mathbb{P}^1(\mathbb{Q}_p) \longrightarrow R \mid \phi \text{ locally constant} \right\} / \{\text{constant}\}$$

There is a natural isomorphism of 1-dim vector spaces:

$$H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p))^+[f] \xrightarrow{\cong} H^1(\Gamma_0(N), \mathbb{Q}_p)^+[f]$$

Flat base change holds:

$$\begin{aligned} & H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Z}_p}(\mathrm{St}_p(\mathbb{Z}_p), \mathbb{Z}_p)) \otimes \mathbb{Q}_p \\ & \xrightarrow{\cong} H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathbb{Q}_p}(\mathrm{St}_p(\mathbb{Q}_p), \mathbb{Q}_p)) \end{aligned}$$

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Definition (Darmon 01)

$$c_{\log_p} = \quad \cdot c_{\text{ord}_p}$$

Step 3: P-adic integration

Remember $\log_p(q_E) = \mathcal{L}_p(E) \cdot \text{ord}_p(q_E)$

Breuil:

$$\rightsquigarrow \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \text{Ext}^1(\mathbb{Q}_p, \text{St}_p^{\text{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\rightsquigarrow c_\lambda: H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)^+[f] \xrightarrow{\cup \mathcal{E}(\lambda)} H^2(\Gamma_0^p(N), \mathbb{Q}_p)^+[f]$$

Fact: c_{ord_p} is an isomorphism of 1-dim vector spaces

Definition (Darmon 01)

$$c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$$

Breuil's extensions

$$\mathrm{Hom}_{\mathrm{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \mathrm{Ext}^1(\mathbb{Q}_p, \mathrm{St}_p^{\mathrm{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\tau_\lambda: \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \tau_\lambda)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} \longrightarrow 0 \\
 & & \downarrow & & & & \uparrow \\
 & & \mathrm{St}_p(\mathbb{Q}_p)^{\mathrm{cont}} & & & & \mathbb{Q}_p
 \end{array}$$

Breuil's extensions

$$\mathrm{Hom}_{\mathrm{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \mathrm{Ext}^1(\mathbb{Q}_p, \mathrm{St}_p^{\mathrm{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\tau_\lambda: \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \tau_\lambda)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} \longrightarrow 0 \\
 & & \downarrow & & & & \uparrow \\
 & & \mathrm{St}_p(\mathbb{Q}_p)^{\mathrm{cont}} & & & & \mathbb{Q}_p
 \end{array}$$

Breuil's extensions

$$\mathrm{Hom}_{\mathrm{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \mathrm{Ext}^1(\mathbb{Q}_p, \mathrm{St}_p^{\mathrm{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\tau_\lambda: \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \tau_\lambda)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} \longrightarrow 0 \\
 & & \downarrow & & & & \uparrow \\
 & & \mathrm{St}_p(\mathbb{Q}_p)^{\mathrm{cont}} & & & & \mathbb{Q}_p
 \end{array}$$

Breuil's extensions

$$\mathrm{Hom}_{\mathrm{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) \longrightarrow \mathrm{Ext}^1(\mathbb{Q}_p, \mathrm{St}_p^{\mathrm{cont}}(\mathbb{Q}_p)), \lambda \longmapsto \mathcal{E}(\lambda)$$

$$\tau_\lambda: \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \tau_\lambda)^{\mathrm{cont}} & \longrightarrow & (\mathrm{Ind}_B^G \mathbb{Q}_p)^{\mathrm{cont}} \longrightarrow 0 \\
 & & \downarrow & & & & \uparrow \\
 & & \mathrm{St}_p(\mathbb{Q}_p)^{\mathrm{cont}} & & & & \mathbb{Q}_p
 \end{array}$$

- 1 Motivation: p -adic periods of elliptic curves
- 2 Darmon's L-invariant
- 3 Big principal series and p -adic families

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+[f]$$

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+[f]$$

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff$$

$$H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+[f]$$

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+[f]$$

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+$$

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff$$

$$H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+[f]$$

A reformulation

Remember $c_{\log_p} = \mathcal{L}_p^D(E) \cdot c_{\text{ord}_p}$

$\dim \text{Hom}_{\text{cont}}(\mathbb{Q}_p^*, \mathbb{Q}_p) = 2$ with basis $\{\text{ord}_p, \log_p\}$

$\rightsquigarrow \langle \log_p - \mathcal{L}_p^D(E) \cdot \text{ord}_p \rangle = \ker(\lambda \mapsto c_\lambda)$

Question: How to produce a non-zero class in $\ker(\lambda \mapsto c_\lambda)$?

$$c_\lambda = 0 \iff$$

$$H^1(\Gamma_0^p(N), \mathcal{E}(\lambda)^\vee)_f^+ \xrightarrow{\neq 0} H^1(\Gamma_0^p(N), \text{St}_p(\mathbb{Q}_p)^\vee)_f^+[f]$$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \quad \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \quad \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \quad \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \quad \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \quad \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Big principal series

$$\tau_\lambda: B \longrightarrow \mathrm{GL}_2(\mathbb{Q}_p), \quad \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \longmapsto \begin{pmatrix} 1 & \lambda(a/d) \\ 0 & 1 \end{pmatrix}$$

$$\tau_\lambda \longleftrightarrow \chi_\lambda: \mathbb{Q}_p^* \longrightarrow (\mathbb{Q}_p[\varepsilon]/\varepsilon^2)^* \text{ such that } \chi_\lambda \equiv 1 \pmod{\varepsilon}$$

Idea: replace $\mathbb{Q}_p[\varepsilon]/\varepsilon^2$ by Tate algebra $\mathcal{O} = \mathbb{Q}_p\langle T \rangle$

Want: $\chi_{\mathcal{O}}: \mathbb{Q}_p^* \longrightarrow \mathcal{O}^*$ such that

- $\chi_{\mathcal{O}} \equiv 1 \pmod{(T)}$ and
- $H^1(\Gamma_0^p(N), \mathrm{Hom}_{\mathcal{O}}((\mathrm{Ind}_B^G \chi_{\mathcal{O}})^{\mathrm{an}}, \mathcal{O}))_f^+ \neq 0$

\rightsquigarrow reduce $\chi_{\mathcal{O}}$ modulo (T^2) to get λ with $c_\lambda = 0$

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^I \chi_{\mathcal{O}})^{\text{an}}$, where

$I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^I \chi \mathcal{O})^{\text{an}}$, where
 $I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^I \chi \mathcal{O})^{\text{an}}$, where
 $I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Theorem (Kohlhaase-Schraen 2012)

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^I \chi_{\mathcal{O}})^{\text{an}}$, where

$I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Theorem (Kohlhaase-Schraen 2012)

$$0 \longrightarrow \text{c-ind}_I^G \mathcal{A} \xrightarrow{U_p - \chi_{\mathcal{O}}(p)} \text{c-ind}_I^G \mathcal{A} \longrightarrow (\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}} \longrightarrow 0$$

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^G \chi_{\mathcal{O}})^{\text{an}}$, where
 $I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Theorem (Kohlhaase-Schraen 2012)

$$0 \longrightarrow \text{c-ind}_I^G \mathcal{A} \xrightarrow{U_p - \chi_{\mathcal{O}}(p)} \text{c-ind}_I^G \mathcal{A} \longrightarrow (\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}} \longrightarrow 0$$

Remark: Theorem holds for more general groups

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^I \chi_{\mathcal{O}})^{\text{an}}$, where
 $I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Theorem (Kohlhaase-Schraen 2012)

$$0 \longrightarrow \text{c-ind}_I^G \mathcal{A} \xrightarrow{U_p - \chi_{\mathcal{O}}(p)} \text{c-ind}_I^G \mathcal{A} \longrightarrow (\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}} \longrightarrow 0$$

Remark: Theorem holds for more general groups
 Multiple U_p -operators

Kohlhaase-Schraen resolution

Consider $\mathcal{A} = (\text{Ind}_{B \cap I}^I \chi_{\mathcal{O}})^{\text{an}}$, where
 $I = \{\gamma \in \text{GL}_2(\mathbb{Z}_p) \mid \gamma \text{ upper triangular modulo } p\}$

Theorem (Kohlhaase-Schraen 2012)

$$0 \longrightarrow \text{c-ind}_I^G \mathcal{A} \xrightarrow{U_p - \chi_{\mathcal{O}}(p)} \text{c-ind}_I^G \mathcal{A} \longrightarrow (\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}} \longrightarrow 0$$

Remark: Theorem holds for more general groups
Multiple U_p -operators \rightsquigarrow Koszul resolution

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1
 U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1
 U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1
 U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$

Overconvergent cohomology in families

Choose $\chi_{\mathcal{O}}|_{\mathbb{Z}_p^*}$ as universal character on weight space, i.e.:

$$\chi_{\mathcal{O}}(x)(T) := \exp_p(T \cdot \log_p(x))$$

Overconvergent families à la Ash-Stevens

$\rightsquigarrow H^1(\Gamma_0(N), \mathcal{A}_f^{\vee})^+$ free \mathcal{O} -module of rank 1

U_p operates via eigenvalue $a_p \in \mathcal{O}^*$

$$\chi_{\mathcal{O}}(x)(T) := a_p(T)^{\text{ord}_p(x)} \cdot \exp_p(T \cdot \log_p(x))$$

$$a_p(0) = a_p(f) = 1 \Rightarrow \chi_{\mathcal{O}} \equiv 1 \pmod{T}$$

$$\text{KS resolution} \Rightarrow H^1(\Gamma_0^p(N), \text{Hom}_{\mathcal{O}}((\text{Ind}_B^G \chi_{\mathcal{O}})^{\text{an}}, \mathcal{O}))_f^+ \neq 0$$

$$\chi_{\mathcal{O}}(x) \pmod{T^2} = 1 + T \left(\frac{d}{dt} a_p \Big|_{t=0} \cdot \text{ord}_p(x) + \log_p(x) \right)$$